

Topology of the space of metrics with positive scalar curvature

Boris Botvinnik
University of Oregon, USA

November 11, 2015
Geometric Analysis in Geometry and Topology, 2015
Tokyo University of Science

Notations:

- W is a compact manifold, $\dim W = d$,
- $\mathcal{R}(W)$ is the space of all Riemannian metrics,
- if $\partial W \neq \emptyset$, we assume that a metric $g = h + dt^2$ near ∂W ;
- R_g is the scalar curvature for a metric g ,
- $\mathcal{R}^+(W)$ is the subspace of metrics with $R_g > 0$;
- if $\partial W \neq \emptyset$, and $h \in \mathcal{R}^+(\partial W)$, we denote

$$\mathcal{R}^+(W)_h := \{g \in \mathcal{R}^+(W) \mid g = h + dt^2 \text{ near } \partial W\}.$$

- “psc-metric” = “metric with positive scalar curvature”.

Existence Question:

- For which manifolds $\mathcal{R}^+(W) \neq \emptyset$?

It is well-known that for a closed manifold W ,

Yamabe invariant

$$\mathcal{R}^+(W) \neq \emptyset \iff Y(W) > 0.$$

Assume $\mathcal{R}^+(W) \neq \emptyset$.

More Questions:

- What is the topology of $\mathcal{R}^+(W)$?
- In particular, what are the **homotopy groups** $\pi_k \mathcal{R}^+(W)$?

Let (W, g) be a spin manifold. Then there is a canonical real spinor bundle $\mathcal{S}_g \rightarrow W$ and a Dirac operator D_g acting on the space $L^2(W, \mathcal{S}_g)$.

Theorem. (Lichnerowicz '60)

$D_g^2 = \Delta_g^s + \frac{1}{4}R_g$. In particular, if $R_g > 0$ then D_g is invertible.

For a manifold W , $\dim W = d$, we obtain a map

$$(W, g) \mapsto \frac{D_g}{\sqrt{D_g^2 + 1}} \in \mathbf{Fred}^{d,0},$$

where $\mathbf{Fred}^{d,0}$ is the space of $\mathcal{C}\ell^d$ -linear Fredholm operators.

The space $\mathbf{Fred}^{d,0}$ also classifies the real K -theory, i.e.,

$$\pi_q \mathbf{Fred}^{d,0} = KO_{d+q}$$

It gives the index map

$$\alpha : (W, g) \mapsto \text{ind}(D_g) = [D_g] \in \pi_0 \mathbf{Fred}^{d,0} = KO_d.$$

The index $\text{ind}(D_g)$ does not depend on a metric g .

Magic of Index Theory gives a map:

$$\alpha : \Omega_d^{\text{Spin}} \longrightarrow KO_d.$$

Thus $\alpha(M) := \text{ind}(D_g)$ gives a topological obstruction to admitting a psc-metric.

Theorem. (Gromov-Lawson '80, Stolz '93) Let W be a spin simply connected closed manifold with $d = \dim W \geq 5$. Then $\mathcal{R}^+(W) \neq \emptyset$ if and only if $\alpha(W) = 0$ in KO_d .



Magic of Topology: There are enough examples of psc-manifolds to generate $\text{Ker } \alpha \subset \Omega_d^{\text{Spin}}$, then we use **surgery**.

Surgery. Let W be a closed manifold, and $S^p \times D^{q+1} \subset W$.



We denote by W' the manifold which is the result of the surgery along the sphere S^p :

$$W' = (W \setminus (S^p \times D^{q+1})) \cup_{S^p \times S^q} (D^{p+1} \times S^q).$$



Codimension of this surgery is $q + 1$.





Surgery Lemma. (Gromov-Lawson) Let g be a metric on W with $R_g > 0$, and W' be as above, where the $q+1 \geq 3$. Then there exists a metric g' on W' with $R_{g'} > 0$.




Surgery Lemma. (Gromov-Lawson) Let g be a metric on W with $R_g > 0$, and W' be as above, where the $q+1 \geq 3$. Then there exists a metric g' on W' with $R_{g'} > 0$.



Surgery Lemma. (Gromov-Lawson) Let g be a metric on W with $R_g > 0$, and W' be as above, where the $q+1 \geq 3$. Then there exists a metric g' on W' with $R_{g'} > 0$.



Surgery Lemma. (Gromov-Lawson) Let g be a metric on W with $R_g > 0$, and W' be as above, where the $q + 1 \geq 3$. Then there exists a metric g' on W' with $R_{g'} > 0$.



Conclusion: Let W and W' be simply connected cobordant spin manifolds, $\dim W = \dim W' = d \geq 5$. Then

$$\mathcal{R}^+(W) \neq \emptyset \iff \mathcal{R}^+(W') \neq \emptyset.$$

Assume $\mathcal{R}^+(W) \neq \emptyset$.

Theorem. (Chernysh, Walsh) Let W and W' be simply connected cobordant spin manifolds, $\dim W = \dim W' \geq 5$. Then

$$\mathcal{R}^+(W) \cong \mathcal{R}^+(W').$$

Let $\partial W = \partial W' \neq \emptyset$, and $h \in \mathcal{R}^+(\partial W)$, then

$$\mathcal{R}^+(W)_h \cong \mathcal{R}^+(W')_h.$$


Questions:

- What is the topology of $\mathcal{R}^+(W)$?
- In particular, what are the **homotopy groups** $\pi_k \mathcal{R}^+(W)$?

Example. Let us show that $\mathbf{Z} \subset \pi_0 \mathcal{R}^+(S^7)$.

Let B be a Bott manifold, i.e. B is a simply connected spin manifold, $\dim B = 8$, with $\alpha(B^8) = \hat{A}(B) = 1$.

Let $\bar{B} := B \setminus (D_1^8 \sqcup D_2^8)$:




Thus $\mathbf{Z} \subset \pi_0 \mathcal{R}^+(S^7)$.

Index-difference construction (N.Hitchin):

Let $g_0 \in \mathcal{R}^+(W) \neq \emptyset$ be a base point, and $g \in \mathcal{R}^+(W)$.

Let $g_t = (1-t)g_0 + tg$.



Fact: The space $(\mathbf{Fred}^{d,0})^+$ is contractible.

The index-difference map: $A_{g_0} : \mathcal{R}^+(W) \longrightarrow \Omega \mathbf{Fred}^{d,0}$.


We obtain a homomorphism:

$$A_{g_0} : \pi_k \mathcal{R}^+(W) \longrightarrow \pi_k \Omega \mathbf{Fred}^{d,0} = KO_{k+d+1}$$

Index-difference construction (N.Hitchin):

Let $g_0 \in \mathcal{R}^+(W) \neq \emptyset$ be a base point, and $g \in \mathcal{R}^+(W)$.

Let $g_t = (1 - t)g_0 + tg$.



Fact: The space $(\mathbf{Fred}^{d,0})^+$ is contractible.

The index-difference map: $A_{g_0} : \mathcal{R}^+(W) \longrightarrow \Omega \mathbf{Fred}^{d,0}$.


We obtain a homomorphism:

$$A_{g_0} : \pi_k \mathcal{R}^+(W) \longrightarrow \pi_k \Omega \mathbf{Fred}^{d,0} = KO_{k+d+1}$$

Index-difference construction (N.Hitchin):

Let $g_0 \in \mathcal{R}^+(W) \neq \emptyset$ be a base point, and $g \in \mathcal{R}^+(W)$.

Let $g_t = (1-t)g_0 + tg$.



Fact: The space $(\mathbf{Fred}^{d,0})^+$ is contractible.

The index-difference map: $A_{g_0} : \mathcal{R}^+(W) \longrightarrow \Omega \mathbf{Fred}^{d,0}$.

We obtain a homomorphism:

$$A_{g_0} : \pi_k \mathcal{R}^+(W) \longrightarrow \pi_k \Omega \mathbf{Fred}^{d,0} = KO_{k+d+1}$$

The classifying space $\mathbf{BDiff}^\partial(W)$. Let W be a connected spin manifold with boundary $\partial W \neq \emptyset$. Fix a collar $\partial W \times (-\varepsilon_0, 0] \hookrightarrow W$. Let

$$\mathrm{Diff}^\partial(W) := \{\varphi \in \mathrm{Diff}(W) \mid \varphi = \mathrm{id} \text{ near } \partial W\}.$$

We fix an embedding $\iota^\partial : \partial W \times (-\varepsilon_0, 0] \hookrightarrow \mathbf{R}^m$ and consider the space of embeddings

$$\mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty}) = \{\iota : W \hookrightarrow \mathbf{R}^{m+\infty} \mid \iota|_{\partial W \times (-\varepsilon_0, 0]} = \iota^\partial\}$$

The group $\mathrm{Diff}^\partial(W)$ acts freely on $\mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty})$ by re-parametrization: $(\varphi, \iota) \mapsto (W \xrightarrow{\varphi} W \xrightarrow{\iota} \mathbf{R}^{m+\infty})$. Then

$$\mathbf{BDiff}^\partial(W) = \mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty}) / \mathrm{Diff}^\partial(W).$$

The space $\mathbf{BDiff}^\partial(W)$ classifies smooth fibre bundles with the fibre W .

The classifying space $\mathbf{BDiff}^\partial(W)$. Let W be a connected spin manifold with boundary $\partial W \neq \emptyset$. Fix a collar $\partial W \times (-\varepsilon_0, 0] \hookrightarrow W$. Let

$$\mathrm{Diff}^\partial(W) := \{\varphi \in \mathrm{Diff}(W) \mid \varphi = \mathrm{id} \text{ near } \partial W\}.$$

We fix an embedding $\iota^\partial : \partial W \times (-\varepsilon_0, 0] \hookrightarrow \mathbf{R}^m$ and consider the space of embeddings

$$\mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty}) = \{\iota : W \hookrightarrow \mathbf{R}^{m+\infty} \mid \iota|_{\partial W \times (-\varepsilon_0, 0]} = \iota^\partial\}$$

The group $\mathrm{Diff}^\partial(W)$ acts freely on $\mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty})$ by re-parametrization: $(\varphi, \iota) \mapsto (W \xrightarrow{\varphi} W \xrightarrow{\iota} \mathbf{R}^{m+\infty})$.

$$\begin{array}{ccc} E(W) & & \\ \downarrow W & & E(W) = \mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty}) \times_{\mathrm{Diff}^\partial(W)} W \\ \mathbf{BDiff}^\partial(W) & & \end{array}$$

The classifying space $\mathbf{BDiff}^\partial(W)$. Let W be a connected spin manifold with boundary $\partial W \neq \emptyset$. Fix a collar $\partial W \times (-\varepsilon_0, 0] \hookrightarrow W$. Let

$$\mathrm{Diff}^\partial(W) := \{\varphi \in \mathrm{Diff}(W) \mid \varphi = \mathrm{id} \text{ near } \partial W\}.$$

We fix an embedding $\iota^\partial : \partial W \times (-\varepsilon_0, 0] \hookrightarrow \mathbf{R}^m$ and consider the space of embeddings

$$\mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty}) = \{\iota : W \hookrightarrow \mathbf{R}^{m+\infty} \mid \iota|_{\partial W \times (-\varepsilon_0, 0]} = \iota^\partial\}$$

The group $\mathrm{Diff}^\partial(W)$ acts freely on $\mathbf{Emb}^\partial(W, \mathbf{R}^{m+\infty})$ by re-parametrization: $(\varphi, \iota) \mapsto (W \xrightarrow{\varphi} W \xrightarrow{\iota} \mathbf{R}^{m+\infty})$.

$$\begin{array}{ccc} E & \xrightarrow{\hat{f}} & E(W) \\ \downarrow W & & \downarrow W \\ B & \xrightarrow{f} & \mathbf{BDiff}^\partial(W) \end{array}$$

Moduli spaces of metrics. Let W be a connected spin manifold with boundary $\partial W \neq \emptyset$, $h_0 \in \mathcal{R}^+(\partial W)$. Recall:

$$\mathcal{R}(W)_{h_0} := \{g \in \mathcal{R}(W) \mid g = h_0 + dt^2 \text{ near } \partial W\},$$

$$\text{Diff}^\partial(W) := \{\varphi \in \text{Diff}(W) \mid \varphi = \text{Id} \text{ near } \partial W\}.$$

The group $\text{Diff}^\partial(W)$ acts freely on $\mathcal{R}(W)_{h_0}$ and $\mathcal{R}^+(W)_{h_0}$:

$$\mathcal{M}(W)_{h_0} = \mathcal{R}(W)_{h_0} / \text{Diff}^\partial(W) = \mathbf{B}\text{Diff}^\partial(W),$$

$$\mathcal{M}^+(W)_{h_0} = \mathcal{R}^+(W)_{h_0} / \text{Diff}^\partial(W).$$

Consider the map $\mathcal{M}^+(W)_{h_0} \rightarrow \mathbf{B}\text{Diff}^\partial(W)$ as a fibre bundle:

$$\mathcal{R}^+(W)_{h_0} \rightarrow \mathcal{M}^+(W)_{h_0} \rightarrow \mathbf{B}\text{Diff}^\partial(W)$$

Let $g_0 \in \mathcal{R}^+(W)_{h_0}$ be a “base point”.


We have the fibre bundle:

$$\begin{array}{c} \mathcal{M}^+(W)_{h_0} \\ \downarrow \mathcal{R}^+(W)_{h_0} \\ \mathbf{BDiff}^\partial(W) \end{array}$$


Let $\varphi : I \rightarrow \mathbf{BDiff}^\partial(W)$ be a loop with $\varphi(0) = \varphi(1) = \mathbf{g}_0$, and

$\tilde{\varphi} : I \rightarrow \mathcal{M}^+(W)_{h_0}$ its lift.

We obtain:



$$\Omega \mathbf{BDiff}^\partial(W) \xrightarrow{e} \mathcal{R}^+(W)_{h_0} \xrightarrow{\text{ind}_{g_0}} \Omega \mathbf{Fred}^{d,0}$$



Let W be a spin manifold, $\dim W = d$. Consider again the index-difference map:

$$\mathbf{ind}_{g_0} : \mathcal{R}^+(W) \longrightarrow \Omega\mathbf{Fred}^{d,0},$$

where $g_0 \in \mathcal{R}^+(W)$ is a “base-point”. In the homotopy groups:

$$(\mathbf{ind}_{g_0})_* : \pi_k \mathcal{R}^+(W) \longrightarrow \pi_k \Omega\mathbf{Fred}^{d,0} = KO_{k+d+1}.$$

Theorem. (BB, J. Ebert, O.Randal-Williams '14) Let W be a spin manifold with $\dim W = d \geq 6$ and $g_0 \in \mathcal{R}^+(W)$. Then


$$\pi_k \mathcal{R}^+(W) \xrightarrow{(\mathbf{ind}_{g_0})_*} KO_{k+d+1} = \begin{cases} \mathbf{Z} & k + d + 1 \equiv 0, 4 \pmod{8} \\ \mathbf{Z}_2 & k + d + 1 \equiv 1, 2 \pmod{8} \\ 0 & \text{else} \end{cases}$$

is non-zero whenever the target group is non-zero.

Remark. This extends and includes results by Hitchin ('75), by Crowley-Schick ('12), by Hanke-Schick-Steimle ('13).



Let $\dim W = d = 2n$. Assume W is a manifold with boundary $\partial W \neq \emptyset$, and W' is the result of an admissible surgery on W . For example:




Then we have:

$$\mathcal{R}^+(D^{2n})_{h_0} \cong \mathcal{R}^+(W')_{h_0},$$

where h_0 is the round metric on S^{2n-1} .

Observation: It is enough to prove the result for $\mathcal{R}^+(D^{2n})_{h_0}$ or any manifold obtained by admissible surgeries from D^{2n} .

We need a particular sequence of **surgeries**:




Here $V_0 = (S^n \times S^n) \setminus D^{2n}$,

$V_1 = (S^n \times S^n) \setminus (D_-^{2n} \sqcup D_+^{2n}), \dots, V_k = (S^n \times S^n) \setminus (D_-^{2n} \sqcup D_+^{2n})$.

Then $W_k := V_0 \cup V_1 \cup \dots \cup V_k = \#_k(S^n \times S^n) \setminus D^{2n}$.

We choose psc-metrics g_j on each V_j which gives the standard round metric h_0 on the boundary spheres.

We need a particular sequence of **surgeries**:




Here $V_0 = (S^n \times S^n) \setminus D^{2n}$,

$V_1 = (S^n \times S^n) \setminus (D_-^{2n} \sqcup D_+^{2n}), \dots, V_k = (S^n \times S^n) \setminus (D_-^{2n} \sqcup D_+^{2n})$.

Then $W_k := V_0 \cup V_1 \cup \dots \cup V_k = \#_k(S^n \times S^n) \setminus D^{2n}$.

We choose psc-metrics g_j on each V_j which gives the standard round metric h_0 on the boundary spheres.




We have the **composition map**

$$\mathcal{R}^+(W_{k-1})_{h_0} \times \mathcal{R}^+(V_k)_{h_0, h_0} \longrightarrow \mathcal{R}^+(W_k)_{h_0}.$$

Gluing metrics along the boundary gives the map:

$$m : \mathcal{R}^+(W_{k-1})_{h_0} \longrightarrow \mathcal{R}^+(W_k)_{h_0}, \quad g \mapsto g \cup g_k$$




We have the **composition map**

$$\mathcal{R}^+(W_{k-1})_{h_0} \times \mathcal{R}^+(V_k)_{h_0, h_0} \longrightarrow \mathcal{R}^+(W_k)_{h_0}.$$

Gluing metrics along the boundary gives the map:

$$m : \mathcal{R}^+(W_{k-1})_{h_0} \longrightarrow \mathcal{R}^+(W_k)_{h_0}, \quad g \mapsto g \cup g_k$$

Magic of Geometry: The map $m : \mathcal{R}^+(W_{k-1})_{h_0} \longrightarrow \mathcal{R}^+(W_k)_{h_0}$ is homotopy equivalence.




Let $s : W_k \hookrightarrow W_{k+1}$ be the inclusion.

It induces the stabilization maps

$$\text{Diff}^\partial(W_0) \rightarrow \cdots \rightarrow \text{Diff}^\partial(W_k) \rightarrow \text{Diff}^\partial(W_{k+1}) \rightarrow \cdots$$

$$\mathbf{BDiff}^\partial(W_0) \rightarrow \cdots \rightarrow \mathbf{BDiff}^\partial(W_k) \rightarrow \mathbf{BDiff}^\partial(W_{k+1}) \rightarrow \cdots$$




Let $s : W_k \hookrightarrow W_{k+1}$ be the inclusion.

It induces the stabilization maps

$$\text{Diff}^\partial(W_0) \rightarrow \cdots \rightarrow \text{Diff}^\partial(W_k) \rightarrow \text{Diff}^\partial(W_{k+1}) \rightarrow \cdots$$

$$\mathbf{BDiff}^\partial(W_0) \rightarrow \cdots \rightarrow \mathbf{BDiff}^\partial(W_k) \rightarrow \mathbf{BDiff}^\partial(W_{k+1}) \rightarrow \cdots$$

Topology-Geometry Magic: the space $\mathbf{BDiff}^\partial(W_k)$ is the moduli space of all Riemannian metrics on W_k which restrict to $h_0 + dt^2$ near the boundary ∂W_k .



Let $s : W_k \hookrightarrow W_{k+1}$ be the inclusion.
 It gives the fiber bundles:

$$\begin{array}{ccccccc}
 \mathcal{M}^+(W_0)_{h_0} & \rightarrow & \mathcal{M}^+(W_1)_{h_0, h_0} & \rightarrow & \cdots & \rightarrow & \mathcal{M}^+(W_k)_{h_0, h_0} & \rightarrow & \cdots \\
 \mathcal{R}^+(W_0)_{h_0} \downarrow & \xrightarrow{\cong} & \mathcal{R}^+(W_0)_{h_0, h_0} \downarrow & \xrightarrow{\cong} & \mathcal{R}^+(W_k)_{h_0, h_0} \downarrow & \xrightarrow{\cong} & & & \\
 \mathbf{BDiff}^\partial(W_0) & \rightarrow & \mathbf{BDiff}^\partial(W_1) & \rightarrow & \cdots & \rightarrow & \mathbf{BDiff}^\partial(W_k) & \rightarrow & \cdots
 \end{array}$$

with homotopy equivalent fibers $\mathcal{R}^+(W_0)_{h_0} \cong \cdots \cong \mathcal{R}^+(W_k)_{h_0, h_0}$

We take a limit to get a fiber bundle:

$$\begin{array}{c} \mathbf{M}_\infty^+ \\ \mathbf{R}_\infty^+ \downarrow \\ \mathbf{B}_\infty \end{array} = \lim_{k \rightarrow \infty} \left(\begin{array}{c} \mathcal{M}^+(W_k)_{h_0, h_0} \\ \mathcal{R}^+(W_k)_{h_0, h_0} \downarrow \\ \mathbf{BDiff}^\partial(W_0) \end{array} \right)$$

where \mathbf{R}_∞^+ is a space homotopy equivalent to $\mathcal{R}^+(W_k)_{h_0, h_0}$.

Remark. We still have the map:

$$\Omega \mathbf{B}_\infty \xrightarrow{e} \mathbf{R}_\infty^+ \xrightarrow{\text{ind}} \Omega \mathbf{Fred}^{d,0}$$

which is consistent with the maps

$$\Omega \mathbf{BDiff}^\partial(W_k) \xrightarrow{e} \mathcal{R}^+(W_k)_{h_0, h_0} \xrightarrow{\text{ind}_{g_0}} \Omega \mathbf{Fred}^{d,0}$$

Magic of Topology: the limiting space $\mathbf{B}_\infty := \lim_{k \rightarrow \infty} \mathbf{BDiff}^\partial(W_k)$ has been understood.

About 10 years ago, **Ib Madsen, Michael Weiss** introduced new technique, parametrized surgery, which allows to describe various

Moduli Spaces of Manifolds.

Theorem. (S. Galatius, O. Randal-Williams) There is a map

$$\mathbf{B}_\infty \xrightarrow{\eta} \Omega_0^\infty \text{MT}\theta_n$$

inducing isomorphism in homology groups.

This gives the fibre bundles:

$$\begin{array}{ccc} \mathbf{M}_\infty^+ & \longrightarrow & \hat{\mathbf{M}}_\infty^+ \\ \mathbf{R}_\infty^+ \downarrow & & \mathbf{R}_\infty^+ \downarrow \\ \mathbf{B}_\infty & \xrightarrow{\eta} & \Omega_0^\infty \text{MT}\Theta_n \end{array}$$

Again, it gives a holonomy map

$$\mathbf{e} : \Omega_0^\infty \text{MT}\Theta_n \longrightarrow \mathbf{R}_\infty^+$$

The space $\Omega_0^\infty \text{MT}\Theta_n$ is the **moduli space of $(n-1)$ -connected $2n$ -dimensional manifolds**.

In particular, there is a map (spin orientation)

$$\hat{\alpha} : \Omega_0^\infty \text{MT}\Theta_n \longrightarrow \mathbf{Fred}^{2n,0}$$

sending a manifold W to the corresponding Dirac operator.

$$\begin{array}{ccc} \Omega\Omega_0^\infty \text{MT}\Theta_n & \xrightarrow{\Omega\hat{\alpha}} & \Omega\mathbf{Fred}^{2n,0} \\ & \searrow e & \nearrow \text{ind} \\ & \mathbf{R}_\infty^+ & \end{array}$$

The space $\Omega_0^\infty \text{MT}\Theta_n$ is the **moduli space of $(n-1)$ -connected $2n$ -dimensional manifolds**.

In particular, there is a map (spin orientation)

$$\hat{\alpha} : \Omega_0^\infty \text{MT}\Theta_n \longrightarrow \mathbf{Fred}^{2n,0}$$

sending a manifold W to the corresponding Dirac operator.

$$\begin{array}{ccc}
 \Omega \Omega_0^\infty \text{MT}\Theta_n & \xrightarrow{\Omega \hat{\alpha}} & \Omega \mathbf{Fred}^{2n,0} \\
 \searrow e & & \nearrow \text{ind} \\
 & \mathbf{R}_\infty^+ &
 \end{array}$$


$$\Omega \mathbf{BDiff}^{\partial}(W) \xrightarrow{e} \mathcal{R}^+(W)_{h_0} \xrightarrow{\text{ind}_{g_0}} \Omega \mathbf{Fred}^{d,0}$$

The space $\Omega_0^\infty \text{MT}\Theta_n$ is the **moduli space of $(n-1)$ -connected $2n$ -dimensional manifolds**.

In particular, there is a map (spin orientation)

$$\hat{\alpha} : \Omega_0^\infty \text{MT}\Theta_n \longrightarrow \mathbf{Fred}^{2n,0}$$

sending a manifold W to the corresponding Dirac operator.



Then we use algebraic topology to compute the homomorphism

$$(\Omega \hat{\alpha})_* : \pi_k(\Omega \Omega_0^\infty \text{MT}\Theta_n) \longrightarrow \pi_k(\Omega \mathbf{Fred}^{2n,0}) = KO_{k+2n+1}$$

to show that it is nontrivial when the target group is non-trivial.



THANK YOU!