

LIST OF PUBLICATIONS
2006-2010

220. Shimizu, M., Shimamura, M., Owaki, T., Asakawa, M., Fujita, K., Kudo, M., Iwakura, Y., Takeda, Y., Mizuguchi, J., and Yoshimoto, T. Antiangiogenic and antitumor activities of IL-27. *J. Immunol.*, **176**, 7317-7324 (2006).
221. Honore, P., Wade, C. L., Zhong, C., Harris, R. R., Wu, C., Ghayur, T., Iwakura, Y., Decker, M. W., Faltynek, C., Sullivan, J., Jarvis, M. F. Interleukin- $\alpha\beta$ gene-deficient mice show reduced nociceptive sensitivity in inflammatory and neuropathic pain but not post-operative pain. *Behav. Brain Res.*, **167**, 355-364 (2006).
222. Kotani, M., Hirata, K., Ogawa, S., Habiro, K., Araki, M., Ishi, T., Ishida, Y., Tanuma, S., Tanabe, K., Toma, Horai, R., Iwakura, Y., and Abe, R. Presence of CD28-dependent and independent pathways in autoimmune arthritis developed in interleukin-1 receptor antagonist-deficient mice. *Arth. Rheum.*, **54**, 473-481 (2006).
223. Deng, X., Yu, Z., Funayama, H., Shoji, N., Sasano, T., Iwakura, Y., Sugawara, S., and Endo, Y. Mutual augmentation of the induction of the histamine-forming enzyme, histidine decarboxylase, between alendronate and immuno-stimulants (IL-1, TNF, and LPS), and its prevention by clodronate. *Toxicol. Appl. Pharmacol.*, **213**, 64-73 (2006).
224. Hadano, S., Benn, S., Kakuta, S., Otomo, A., Sudo, K., Kunita, R., Suzuki-Utsunomiya, K., Mizumura, H., Shefner, J. M., Cox, G. A., Iwakura, Y., Brown, R. H. Jr., and Ikeda, J. Loss of ALS2 in mice produces mild, age-dependent loss of cerebellar Purkinje cells and subtle pathology of large spinal motor neurons without an obvious motor phenotype. *Human Molecular Genetics*, **15**, 233-250 (2006).
225. Matsuki, T., Nakae, S., Sudo, K., Horai, R., and Iwakura, Y. Abnormal T cell activation caused by the imbalance of the IL-1/IL-1 receptor antagonist system is responsible for the development of

- experimental autoimmune encephalomyelitis. *Int. Immunol.*, **18**, 399-407 (2006).
226. Ishigame, H., Nakajima, A., Saijo, S., Komiyama, Y., Mastuki, T., Nakae, S., Horai, R., Kakuta, S., and Iwakura, Y. The role of TNF α and IL-17 in the development of excess IL-1 signaling-induced inflammatory diseases in IL-1 receptor antagonist-deficient mice. *Ernst Schering Res. Found. Workshop*, **56**, 129-153 (2006).
227. Zhu, Y., Saito, K., Murakami, Y., Asano, M., Iwakura, Y., and Seishima^M. Early increase in mRNA levels of pro-inflammatory cytokines and their interactions in the mouse hippocampus after transient global ischemia. *Neuroscience Letter*, **393**, 122-126 (2006).
228. Vonk, A. G., Netea, M. G., van Krieken, J. H., Iwakura, Y., van der Meer, J. W. M., and Kullberg, J. B. Endogenous interleukin-1 α and interleukin-1 β are crucial for host defense against disseminated candidiasis. *J. Infect. Dis.*, **193**, 1419-1426 (2006).
229. Chida, D., Osaka, T., Hashimoto, O., and Iwakura, Y. Combined IL-6 and IL-1 deficiency causes obesity in young mice. *Diabetes*, **55**, 971-977 (2006).
230. Nambu, A., Nakae, S., and Iwakura, Y. IL-1 β , but not IL-1 α , is required for antigen-specific T cell activation and the induction of local inflammation in the delayed-type hypersensitivity responses. *Int. Immunol.*, **18**, 701-712 (2006).
231. Sasaki, S., Tagawa, Y., Iwakura, Y., and Nakane, A. The role of gamma interferon in acquired host resistance against *Staphylococcus aureus* infection in mice. *FEMS Immunol. Med. Microbiol.*, **46**, 367-374 (2006).
232. Tateno, K., Minamino, T., Toko, H., Akazawa, H., Shimizu, N., Takeda, S., Kunieda, T., Miyauchi, H., Oyama, T., Matsuura, K., Nishi, J. I., Kobayashi, Y., Nagai, T., Kuwabara, Y., Iwakura, Y., Nomura, F., Saito, Y., and Komuro, I. Critical roles of

- muscle-secreted angiogenic factors in therapeutic neovascularization. *Circ. Res.*, **98**, 1194-1202 (2006).
233. O'Sullivan, B. J., Thomas, H. E., Pai, S., Santamaria, P., Iwakura, Y., Steptoe, R. J., Kay, T. W. H., and Thomas, R. IL-1 β breaks tolerance through expansion of CD25⁺ effector T cells. *J. Immunol.*, **176**, 7278-7287 (2006).
 234. Iwakura, Y., and Ishigame, H. The IL-23/IL-17 axis in inflammation. *J. Clin. Invest.*, **116**, 1218-1222 (2006).
 235. Ohtaki, H., Nakamachi, T., Dohi, K., Aizawa, Y., Takaki, A., Hodoyama, K., Yofu, S., Hashimoto, H., Shintani, N., Baba, A., Kopf, M., Iwakura, Y., Matsuda, K., Arimura, A., and Shioda, S. Pituitary adenylate cyclase-activating polypeptide (PACAP) decreases ischemic neuronal cell death in association with interleukin-6. *Proc. Natl. Acad. Sci. USA*, **103**, 7488-7493 (2006).
 236. Komiyama, Y., Nakae, S., Matsuki, T., Nambu, A., Ishigame, H., Kakuta, S., Sudo, K., and Iwakura, Y. IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J. Immunol.*, **177**, 566-573 (2006).
 237. Uetani, N., Chagnon, M. J., Kennedy, T. E., Iwakura, Y., and Tremblay, M. L. Mammalian motoneuron axon targeting requires receptor protein tyrosine phosphatases sigma and delta. *J. Neurosci.*, **26**, 5872-5880 (2006).
 238. Nakazawa, T., Komai, S., Watabe, A. M., Kiyama, Y., Fukaya, M., Arima-Yoshida, F., Horai, R., Sudo, K., Ebine, K., Delawary, M., Goto, J., Umemori, H., Tezuka, T., Iwakura, Y., Watanabe, M., Yamamoto, T., and Manabe, T. NR2B tyrosine phosphorylation modulates fear learning as well as amygdaloid synaptic plasticity. *EMBO J.*, **25**, 2867-2877 (2006).
 239. Murakami, M., Iwai, S., Hiratsuka, S., Yamauchi, M., Nakamura, K., Iwakura, Y., and Shibuya, M. Signaling of vascular endothelial growth factor receptor-1 tyrosine kinase promotes rheumatoid

- arthritis through activation of monocyte/macrophage. *Blood*, **108**, 1849-1856 (2006).
240. Okada, K., Inoue, A., Okada, M., Murata, Y., Kakuta, S., Jigami, T., Kubo, S., Shiraishi, H., Eguchi, K., Motomura, M., Akiyama, T., Iwakura, Y., Higuchi, O., and Yamanashi, Y. The muscle protein Dok-7 is essential for neuromuscular synaptogenesis. *Science*, **312**, 1802-1805 (2006).
 241. Fujikado, N., Saijo, S., and Iwakura, Y. Identification of arthritis-related gene clusters by microarray analysis of two independent mouse models for rheumatoid arthritis. *Arthritis Res. Ther.*, **8**, R100, 1-13 (2006).
 242. Furuta, T., Kikuchi, T., Iwakura, Y., and Watanabe, N. Protective roles of mast cells and mast cell-derived tumor necrosis factor in murine malaria. *J. Immunol.*, **177**, 3294-302 (2006).
 243. Voronov, E., Dayan, M., Zinger, H., Gayvoronsky, L., Lin, J. P., Iwakura, Y., Apte, R. N., and Mozes, E. IL-1beta-deficient mice are resistant to induction of experimental SLE. *Eur. Cytokine Netw.*, **17**, 109-16 (2006).
 244. Furuichi, K., Wada, T., Iwata, Y., Kokubo, S., Hara, A., Yamahana, J., Sugaya, T., Iwakura, Y., Matsushima, K., Asano, M., Yokoyama, H., and Kaneko, S. Interleukin-1-dependent sequential chemokine expression and inflammatory cell infiltration in ischemia-reperfusion injury. *Crit. Care Med.*, **34**, 2447-2455 (2006).
 245. Konishi, Y., Ikeda, K., Iwakura, Y., and Kawakami, K. Six1 and Six4 promote survival of sensory neurons during early trigeminal gangliogenesis. *Brain Res.*, **1116**, 93-102 (2006).
 246. Maedler, K., Schumann, D. M., Sauter, N., Ellingsgaard, H., Bosco, D., Baertshiger, R., Iwakura, Y., Oberholzer, J., Wollheim, C. B., Gauthier, B. R., and Donath, M. Y. Low concentration of interleukin-1 β induces FLICE-inhibitory protein-mediated β -cell proliferation in human pancreatic islets. *Diabetes*, **55**, 2713-2722 (2006).

247. Sato, K., Suematsu, A., Okamoto, K., Yamaguchi, A., Morishita, Y., Kadono, Y., Tanaka, S., Kodama, T., Shizuo, A., Iwakura, Y., Cua, D. J., and Takayanagi, H. T_H17 functions as an osteoclastogenic helper T cell subset that links T cell activation and bone destruction. *J. Exp. Med.*, **203**, 2673-2682 (2006).
248. Ito, R., Shin-Ya, M., Kishida, T., Urano, A., Takada, R., Sakagami, J., Imanishi, J., Kita, M., Ueda, Y., Iwakura, Y., Kataoka, K., Okanoue, T., and Mazda, O. Interferon-gamma is causatively involved in experimental inflammatory bowel disease in mice. *Clin. Exp. Immunol.*, **146**, 330-338 (2006).
249. Tsurutani, N., Yasuda, J., Yamamoto, N., Choi, B., Kadoki, M., and Iwakura, Y. Nuclear import of the pre-integration complex is blocked upon infection by HIV-1 in mouse cells. *J. Virol.*, **81**, 677-688 (2007).
250. Saijo, S., Fujikado, N., Furuta, T., Chung, S., Kotaki, H., Seki, K., Sudo, K., Akira, S., Adachi, Y., Ohno, N., Kinjo, T., Nakamura, K., Kawakami, K., and Iwakura, Y. Dectin-1 is required for host defense against *Pneumocystis carinii* but not against *Candida albicans*. *Nature Immunol.*, **8**, 39-46 (2007).
251. Hirota, K., Hashimoto, M., Yoshitomi, H., Tanaka, S., Nomura, T., Yamaguchi, T., Iwakura, Y., Sakaguchi, N., and Sakaguchi, S. T cell self-reactivity forms cytokine milieu for spontaneous development of IL-17⁺ helper T cells that cause autoimmune arthritis. *J. Exp. Med.*, **204**, 41-47 (2007).
252. Nigrovic, P. A., Binstadt, B. A., Monach, P. A., Johnsen, A., Gurish, M., Iwakura, Y., Benoist, C., Mathis, D., and Lee, D. M. Mast cells contribute to initiation of autoantibody-mediated arthritis via IL-1. *Proc. Natl. Acad. Sci. USA*, **104**, 2325-2330, (2007).
253. Nakamatsu, M., Yamamoto, N., Nakasone, C., Kinjo, T., Miyagi, K., Uezu, K., Nakamura, K., Nakayama, T., Taniguchi, M., Iwakura, Y., Kaku, M., Fujita, J., and Kawakami, K. Role of interferon-gamma in $V\alpha14^+$ natural killer T cell-mediated host defense against

- Streptococcus pneumoniae infection in murine lungs. *Microbes Infect.*, **9**, 364-374 (2007).
254. Kamari, Y., Werman-Venkert, R., Shaish, A., Werman, A., Harari, A., Gonen, A., Voronov, E., Grosskopf, I., Sharabi, Y., Grossman, E., Iwakura, Y., Dinarello, C. A., Apte, R. N., and Harats, D. Differential role and tissue specificity of interleukin-1alpha gene expression in atherogenesis and lipid metabolism. *Atherosclerosis*, **195**, 31-38 (2007).
 255. Sugihara, T., Sekine, C., Nakae, T., Kohyama, K., Harigai, M., Iwakura, Y., Matsumoto, Y., Miyasaka, N., and Kohsaka, H. A new murine model defines critical mediators in the pathology and treatment of polymyositis. *Arthritis Rheum.*, **56**, 1304-1314 (2007).
 256. Adachi, K., Soeta-Saneyoshi, C., Sagara, H., and Iwakura, Y. A crucial role of *Bysl* in mammalian preimplantation development as an integral factor for 40S ribosome biogenesis. *Mol. Cell. Biol.*, **27**, 2202-2214 (2007).
 257. Nakae, S., Iwakura, Y., Suto, H., and Galli, S. J. Phenotypic differences between Th1 and Th17 cells and negative regulation of Th1 cell-differentiation by IL-17. *J. Leu. Biol.*, **81**, 1258-1268 (2007).
 258. Ashino, T., Arima, Y., Shioda, S., Iwakura, Y., Numazawa, S., and Yoshida, T. Effect of interleukin-6 on CYP3A11 and metallothionein-1/2 expression in arthritic mouse liver. *Eur. J. Pharm.*, **558**, 199-207 (2007).
 259. Onodera, S., Ohshima, S., Tohyama, H., Yasuda, K., Nishihira, J., Iwakura, Y., Matsuda, I., Minami, A., and Koyama, Y. A novel DNA vaccine targeting macrophage migration inhibitory factor protects joints from inflammation and destruction in murine models of arthritis. *Arthritis Rheum.*, **56**, 521-530 (2007).
 260. Ishida, M., Choi, J. H., Hirabayashi, K., Matsuwaki, T., Suzuki, M., Yamanouchi, K., Horai, R., Sudo, K., Iwakura, Y., and Nishihara, M. Reproductive phenotypes in mice with targeted disruption of the

- 20 α -hydroxysteroid dehydrogenase gene. *J. Reprod. Dev.*, **53**, 499-508 (2007).
261. Krelín, Y., Voronov, E., Dotan, S., Elkabets, M., Reich, E., Fogel, M., Huszar, M., Iwakura, Y., Segal, S., Dinarello, C. A., and Apte, R. N. Interleukin-1 β -driven inflammation promotes the development and invasiveness of chemical carcinogen-induced tumors. *Cancer Res.*, **67**, 1062-1071 (2007).
262. Ohama, T., Hori, M., Momotani, E., Iwakura, Y., Guo, F., Kishi, H., Kobayashi, S., and Ozaki, H. Intestinal inflammation down-regulates smooth muscle CPI-17 through induction of TNF- α and causes motility disorders. *Am. J. Physiol. Gastrointest. Liver Physiol.*, **292**, G1429-1438 (2007).
263. Umemura, M., Yahagi, A., Hamada, S., Begum, M-D., Watanabe, H., Kawakami, K., Suda, T., Sudo, K., Nakae, S., Iwakura, Y., and Matsuzaki, G. IL-17-mediated regulation of innate and acquired immune response against pulmonary *Mycobacterium bovis bacilli Calmette-Guerin* infection. *J. Immunol.*, **178**, 3786-3796 (2007).
264. Togashi, Y., Chamoto, K., Wakita, D., Tsutsumi, N., Iwakura, Y., Matsubara, N., Kitamura, H., and Nishimura, T. Natural killer T cells from interleukin-4-deficient mice are defective in early interferon-gamma production in response to alpha-galactosylceramide. *Cancer Sci.*, **98**, 721-725 (2007).
265. Hayashi, T., Ishida, Y., Kimura, A., Iwakura, Y., Mukaida, N., and Kondo, T. IFN- γ protects cerulein-induced acute pancreatitis by repressing NF-kB activation. *J. Immunol.*, **178**, 7385-7394 (2007).
266. Kaiga, T., Sato, M., Kaneda, H., Iwakura, Y., Takayama, T., and Tahara, H. Systemic administration of IL-23 induces potent anti-tumor immunity primarily mediated through Th1-type response in association with the endogenously expressed IL-12. *J. Immunol.*, **178**, 7571-7580 (2007).
267. Numasaki, M., Tagawa, M., Iwata, F., Suzuki, T., Nakamura, A., Okada, M., Iwakura, Y., Aiba, S., and Yamaya, M. IL-28 elicits

- antitumor responses against murine fibrosarcoma. *J. Immunol.*, **178**, 5086-5098 (2007).
268. Sato, N., Kinbara, M., Kuroishi, T., Kimura, K., Iwakura, Y., Ohtsu, H., Sugawara, S., and Endo, Y. Lipopolysaccharide promotes and augments metal allergies in mice, dependent on innate immunity and histidine decarboxylase. *Clin. Exp. Allergy*, **37**, 743-751 (2007).
269. Hara, H., Ishihara, C., Takeuchi, A., Imanishi, T., Xue, L., Morris, S. W., Inui, M., Takai, T., Shibuya, A., Saijo, S., Iwakura, Y., Ohno, N., Koseki, H., Yoshida, H., Penninger, J. M., and Saito, T. The adaptor protein CARD9 is essential for the activation of myeloid cells through ITAM-associated and Toll-like receptors. *Nature Immunol.*, **8**, 619-629 (2007).
270. Venkataraman, T., Valdes, M., Elsby, R., Kakuta, S., Caceres, G., Saijo, S., Iwakura, Y., and Barber, G. N. Loss of DExD/H box RNA Helicase LGP2 manifests disparate antiviral responses. *J. Immunol.*, **178**, 6444-6455 (2007).
271. Ozato, K., Uno, K., and Iwakura, Y. Another road to interferon: Yasuichi Nagano's journey. *J. Interferon Cytokine Res.*, **27**, 349-352. (2007).
272. Takagi, K., Matsumura, S., Okuda-Ashitaka, E., Okuda, K., Watanabe, J., Takahashi, H., Iwakura, Y., and Ito, S. Interleukin-1 is not essential for expression of inducible NOS in hepatocytes induced by lipopolysaccharide *in vivo*. *Nitric Oxide*, **16**, 433-441, (2007).
273. Unezaki, S., Horai, R., Sudo, K., Iwakura, Y., and Ito, S. Ovol2/Movo, a homologue of *Drosophila* ovo, is required for angiogenesis, heart formation and placental development in mice. *Genes Cells*, **12**, 773-785 (2007).
274. Kohyama, S., Ohno, S., Isoda, A., Moriya, O., Belladonna, M. L., Hayashi, H., Iwakura, Y., Yoshimoto, T., Akatsuka, T., and Matsui, M. IL-23 enhances host defense against vaccinia virus infection via a mechanism partly involving IL-17. *J. Immunol.*, **179**,

3917-3925 (2007).

275. Kina, S., Tezuka, T., Kusakawa, S., Kishimoto, Y., Kakizawa, S., Hashimoto, K., Ohsugi, M., Kiyama, Y., Horai, R., Sudo, K., Kakuta, S., Iwakura, Y., Iino, M., Kano, M., Manabe, T., and Yamamoto, T. Involvement of protein-tyrosine phosphatase PTPMEG in motor learning and cerebellar long-term depression. *Eur. J. Neurosci.*, **26**, 2269-2278, (2007).
276. Nakatani, Y., Hokonohara, Y., Kakuta, S., Sudo, K., Iwakura, Y., and Kudo, I. Knockout mice lacking cPGES/p23, a constitutively expressed PGE(2) synthetic enzyme, are peri-natally lethal. *Biochem. Biophys. Res. Commun.*, **362**, 387-92, (2007).
277. Kayasuga, Y., Chiba, S., Suzuki, M., Kikusui, T., Matsuwaki, T., Yamanouchi, K., Kotaki, H., Horai, R., Iwakura, Y., and Nishihara, M. Alteration of behavioural phenotype in mice by targeted disruption of the progranulin gene. *Behav. Brain Res.*, **185**, 110-118 (2007).
278. Tamagawa, A., Kolosova, I., Endo, Y., Gerlinskaya, L., Iwakura, Y., and Moshikin, M. Interleukin-1 deficiency and aggressiveness in male mice. In “*Psychoneuroendocrinology Research Trends*”, M. T. Czerbnska ed., Nova Science Publishers, Inc., (ISBN 978-1-60021-665-7), pp1-16 (2007).
279. Kagari, T., Tanaka, D., Doi, H., Iwakura, Y., and Shimozato, T. Anti-type II collagen antibody accelerates arthritis via CXCR2-expressing cells in IL-1 receptor antagonist-deficient mice. *Eur. J. Immunol.*, **37**, 2753-2763 (2007).
280. Chida, D., Nakagawa, S., Nagai, S., Sagara, H., Katsumata, H., Imaki, T., Suzuki, H., Mitani, F., Ogishima, T., Shimizu, C., Kotaki, H., Kakuta, S., Sudo, K., Koike, T., Kubo, M., and Iwakura, Y. Melanocortin receptor 2 is required for adrenal gland development, steroidogenesis and neonatal gluconeogenesis. *Proc. Natl. Acad. Sci. USA*, **104**, 18205-18210 (2007).
281. Chida, D., and Iwakura, Y. Peripheral TNF α , but not peripheral

- IL-1, requires endogenous IL-1 or TNF α induction in the brain for the febrile response. *Biochem. Biophys. Res. Commun.*, **364**, 765-770 (2007).
282. Noguchi, D., Wakita, D., Tajima, M., Ashino, S., Iwakura, Y., Zhang, Y., Chamoto, K., Kitamura, H., and Nishimura, T. Blocking of IL-6 signaling pathway prevents CD4⁺ T cell-mediated colitis in a Th17-independent manner. *Int. Immunol.*, **19**, 1431-1440 (2007).
283. Miller, L. S., Pietras, E. M., Uricchio, L. H., Hirano, K., Rao, S., Lin, H., O'Connell, R. M., Iwakura, Y., Cheung, A. L., Cheng, G., and Modlin, R. L. Inflammasome-mediated production of IL-1 β is required for neutrophil recruitment against *Staphylococcus aureus* in vivo. *J. Immunol.*, **179**, 6933-6942 (2007).
284. Nakamura, K., Kinjo, T., Saijo, S., Miyazato, A., Adachi, Y., Ohno, N., Fujita, J., Kaku, M., Iwakura, Y., and Kawakami, K. Dectin-1 is not required for the host defense to *Cryptococcus neoformans*. *Microbiol. Immunol.*, **51**, 1115-1119 (2007).
285. Okumura, A., Saito, T., Otani, I., Kojima, K., Yamada, Y., Ishida-Okawara, A., Nakazato, K., Asano, M., Kanayama, K., Iwakura, Y., Suzuki, K., and Yamagoe, S. Suppressive role of leukocyte cell-derived chemotaxin 2 in mouse anti-type II collagen antibody-induced arthritis. *Arthritis Rheum.*, **58**, 413-421 (2008).
286. Fujikado, N., Saijo, S., Yonezawa, T., Shimamori, K., Ishii, A., Sugai, S., Kotaki, H., Sudo, K., Nose, M., and Iwakura, Y. Dcir deficiency causes development of autoimmune diseases in mice due to excess expansion of dendritic cells. *Nature Med.*, **14**, 176-180 (2008).
287. Ashino, T., Yamanaka, R., Yamamoto, M., Shimokawa, H., Sekikawa, K., Iwakura, Y., Shioda, S., Numazawa, S., and Yoshida, T. Negative feedback regulation of lipopolysaccharide-induced inducible nitric oxide synthase gene expression by heme oxygenase-1 induction in macrophages. *Mol. Immunol.*, **45**, 2106-2115 (2008).
288. Yoshimura, T., Sonoda, K., Miyazaki, Y., Iwakura, Y., Ishibashi, T., Yoshimura, A., and Yoshida, H. Differential roles for IFN- γ and

- IL-17 in experimental autoimmune uveoretinitis. *Int. Immunol.*, **20**, 209-214 (2008).
289. Honda, A., Abe, R., Makino, T., Norisugi, O., Fujita, Y., Watanabe, H., Nishihira, J., Yamagishi, S., Iwakura, Y., Shimizu, H., and Shimizu, T. Interleukin-1 β and macrophage migration inhibitory factor (MIF) in dermal fibroblasts mediate UVA induced matrix metalloproteinase-1 expression. *J. Dermatol. Sci.*, **49**, 63-72 (2008).
290. Pichavant, M., Goya, S., Meyer, E. H., Johnston, R. A., Kim, H. Y., Matangkasombut, P., Zhu, M., Iwakura, Y., Savage, P. B., DeKruyff, R. H., Shore, S. A., and Umetsu, D. T. Ozone exposure in a mouse model induces airway hyperreactivity that requires the presence of Natural Killer T cells and IL-17. *J. Exp. Med.*, **205**, 385-393 (2008).
291. Ohsugi, M., Adachi, K., Horai, R., Kakuta, S., Sudo, K., Kotaki, H., Tokai-Nishizumi, N., Sagara, H., Iwakura, Y., and Yamamoto, T. Kid-mediated chromosome compaction ensures proper nuclear envelope formation. *Cell*, **132**, 771-782 (2008).
292. Nakamura, K., Miyazato, A., Gang, X., Hatta, M., Inden, K., Aoyagi, T., Takeda, K., Akira, S., Saijo, S., Iwakura, Y., Adachi, Y., Ohno, N., Suzuki, K., Fujita, J., Kaku, M., and Kawakami, K. Deoxynucleic acids from *Cryptococcus neoformans* activate myeloid dendritic cells via a TLR9-dependent pathway. *J. Immunol.*, **180**, 4067-4074 (2008).
293. Aujla, S. J., Fei, M., Pociask, D. A., Reinhart, T. A., Edeal, J., Gaus, K., Kreindler, J. L., Dubin, P. J., Zheng, M., Pilewski, J. M., Myerburg, M. M., Mason, C. A., Iwakura, Y., and Kolls, J. K. IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. *Nature Med.*, **14**, 275-281 (2008).
294. Luger, D., Silver, P. B., Tang, J., Cua, D., Chen, Z., Iwakura, Y., Bowman, E. P., Sgambellone, N. M., Chan, C.-C. and Caspi, R. R. Either a Th17 or a Th1 effector response can drive autoimmunity: conditions of disease induction affect dominant effector category. *J.*

- Exp. Med.*, **205**, 799-810 (2008).
295. Fukushima, A., Sumi, T., Ishida, W., Yamada, J., Iwakura, Y., and Ueno, H. Endogenous IL-17 does not play a significant role in the development of experimental murine allergic conjunctivitis. *Int. Arch. Allergy Imm.*, **147**, 206-212 (2008)
296. Izcue, A., Hue, S., Buonocore, S., Arancibia-Carcamo, C. V., Ahern, P. P., Iwakura, Y., Maloy, K. J., and Powrie, F. Interleukin-23 restrains regulatory T cell activity to drive T cell-dependent colitis. *Immunity*, **28**, 559-570 (2008).
297. Tajima, M., Wakita, D., Noguchi, D., Chamoto, K., Yue, Z., Fugo, K., Ishigame, H., Iwakura, Y., Kitamura, H., and Nishimura, T. IL-6-dependent spontaneous proliferation is required for the induction of colitogenic IL-17-producing CD8⁺ T cells. *J. Exp. Med.*, **205**, 1019-1027 (2008).
298. Sakai, M., Iwakawa, M., Iwakura, Y., Ohta, T., Tsujii, H., and Imai, T. CD44 and Bak expression in IL-6 or TNF- α gene knockout mice after whole lung irradiation. *J. Radiat. Res. (Tokyo)*, **49**, 409-416 (2008)
299. Doodes, P. D., Cao, Y., Hamel, K., Wang, Y., Hutas, G., Iwakura, Y., and Finnegan, A. Development of proteoglycan-induced arthritis is independent of IL-17. *J. Immunol.*, **181**, 329-337 (2008).
300. Bai, Y., Liu, R., Huang, D., La Cava, A., Tang, Y., Iwakura, Y., Campagnolo, D. I., Vollmer, T. L., Ransohoff, R. M., and Shi, F. CCL2 recruitment of IL-6 producing CD11b⁺ monocytes to the draining lymph nodes during the initiation of TH17-dependent B cell-mediated autoimmunity. *Eur. J. Immunol.*, **38**, 1877-1888 (2008).
301. Chida, D., Hashimoto, O., Kuwahara, M., Sagara, H., Osaka, T., Tsubone, H., and Iwakura, Y. Increased fat-to-carbohydrate oxidation ratio in IL-1Ra^{-/-} mice on a high fat diet is associated with increased sympathetic tone. *Diabetologia*, **51**, 1698-1706 (2008).
302. Lees, J. R., Iwakura, Y., and Russell, J. H. Host T cells are the main

producers of IL-17 within the central nervous system during initiation of experimental autoimmune encephalomyelitis induced by adoptive transfer of Th1 cell lines. *J. Immunol.*, **180**, 8066-8072 (2008).

303. Schulz, S. M., Köhler, G., Holscher, C., Iwakura, Y., and Alber, G. IL-17A is produced by Th17 cells, $\gamma\delta$ T cells and other CD4⁺ lymphocytes during infection with *Salmonella enterica* serovar Enteritidis and has a mild effect in bacterial clearance. *Int. Immunol.*, **9**, 1129-1138 (2008).
304. Kafka, D., Ling, E., Feldman, G., Benharroch, D., Voronov, E., Givon-Laci, N., Iwakura, Y., Dagan, R., Apte, R. N., and Mizrahi-Nebenzahi, Y. Contribution of IL-1 to resistance to *Streptococcus pneumoniae* infection. *Int. Immunol.*, **9**, 1139-1146 (2008).
305. Kubota, N., Kubota, T., Itoh, S., Kumagai, H., Kozono, H., Takamoto, I., Mineyama, T., Ogata, H., Tokuyama, K., Ohsugi, M., Sakako, T., Moroi, M., Sugi, K., Kakuta, S., Iwakura, Y., Noda, T., Ohnishi, S., Nagai, R., Tobe, K., Terauchi, Y., Ueki, K., and Kadowaki, T. Dynamic functional relay between insulin receptor substrate 1 and 2 in hepatic insulin signaling during fasting and feeding. *Cell Metab.*, **8**, 49-64 (2008).
306. Nakamura, K., Miyazato, A., Koguchi, Y., Adachi, Y., Ohno, N., Saijo, S., Iwakura, Y., Takeda, K., Akira, S., Fujita, J., Ishii, K., Kaku, M., and Kawakami, K. Toll-like receptor 2 (TLR2) and dectin-1 contribute to the production of IL-12p40 by bone marrow-derived dendritic cells infected with *Penicillium marneffei*. *Microbes Infect.*, **10**, 1223-1227 (2008).
307. Shiomi, S., Toriie, A., Imamura, S., Konishi, H., Mitsufuji, S., Iwakura, Y., Yamaoka, Y., Ota, H., Yamamoto, T., Imanishi, J., and Kita, M. IL-17 is involved in *Helicobacter pylori*-induced gastric inflammatory responses in a mouse model. *Helicobacter*, **13**, 518-524 (2008).
308. Iwakura, Y., Nakae, S., Saijo, S., and Ishigame, H. The roles of

IL-17A in inflammatory immune responses and host defense against pathogens. *Immunological Rev.*, **226**, 57-79 (2008).

309. Hamada, S., Umemura, M., Shiono, T., Tanaka, K., Yahagi, A., Begum, M. D., Oshiro, K., Okamoto, Y., Watanabe, H., Kawakami, K., Roark, C., Born, W. K., O'Brien, R., Ikuta, K., Ishikawa, H., Nakae, S., Iwakura, Y., Ohta, T., and Matsuzaki, G. L-17A produced by gammadelta T cells plays a critical role in innate immunity against *Listeria monocytogenes* infection in the liver. *J. Immunol.*, **181**, 3456-3463 (2008).
310. Ito, R., Kita, M., Shin-Ya, M., Kishida, T., Urano, A., Takada, R., Sakagami, J., Imanishi, J., Iwakura, Y., Okanoue, T., Yoshikawa, T., Kataoka, K., and Mazda, O. Involvement of IL-17A in the pathogenesis of DSS-induced colitis in mice. *Biochem. Biophys. Res. Commun.*, **377**, 12-16 (2008).
311. Murayama, H., Takahashi, M., Takamoto, M., Shiba, Y., Ise, H., Koyama, J., Tagawa, Y., Iwakura, Y., and Ikeda, U. Deficiency of tumor necrosis factor- α and interferon- γ in bone marrow cells synergistically inhibits neointimal formation following vascular injury. *Cardiovascular Res.*, **80**, 175-180 (2008).
312. Wakashin, H., Hirose, K., Maezawa, Y., Kagami, S., Suto, A., Watanabe, N., Saito, Y., Hatano, M., Tokuhisa, T., Iwakura, Y., Puccetti, P., Iwamoto, I., and Nakajima, H. IL-23 and Th17 cells enhance Th2 cell-mediated eosinophilic airway inflammation in mice. *Am. J. Respir. Crit. Care Med.*, **178**, 1023-1032 (2008).
313. Teruya, H., Tomita, M., Senba, M., Ishikawa, C., Tamayose, M., Miyazato, A., Yara, S., Tanaka, Y., Iwakura, Y., Fujita, J., and Mori, N. Human T-cell leukemia virus type I infects human lung epithelial cells and induces gene expression of cytokines, chemokines and cell adhesion molecules. *Retrovirology*, **5**, 86 (2008).
314. Kappel, L. W., Goldberg, G. L., King, C. G., Suh, D., Smith, O. M., Ligh, C., Holland, A., Grubin, J., Liu, C., Iwakura, Y., Mark, N., Heller, G., and van den Brink, M. R. M. IL-17 contributes to

CD4-mediated graft-versus-host disease. *Blood*, **113**, 628-636 (2008).

315. Schulz, S. M., Köhler, G., Schütze, N., Knauer, J., Straubinger, R. K., Chackerian, A. A., Witte, E., Wolk, K., Sabat, R., Iwakura, Y., Holscher, C., Müller, U., Kastelein, R. A., and Alber, G. Protective immunity to systemic infection with *Salmonella enterica* serovar Enteritidis in the absence of IL-12 is associated with IL-23-dependent IL-22 but not IL-17. *J. Immunol.*, **181**, 7891-7901 (2008).
316. Ogura, H., Murakami, M., Okuyama, Y., Tsuruoka, M., Kitabayashi, C., Kanamoto, M., Mishihara, M., Iwakura, Y., and Hirano, T. Interleukin-17 promotes autoimmunity by triggering a positive-feedback loop via Interleukin-6 induction. *Immunity*, **29**, 628-636 (2008).
317. Eigenbrod, T., Park, J-H., Harder, J., Iwakura, Y., and Nunez, G. Critical role for mesothelial cells in necrosis-induced inflammation through the recognition of IL-1 α released from dying cells. *J. Immunol.*, **181**, 8194-8198 (2008).
318. Atsumi, T., Sato, M., Kamimura, D., Moroi, A., Iwakura, Y., Betz, U. A., Yoshimura, A., Nishihara, M., Hirano, T., and Murakami, M. IFN- γ expression in CD8⁺ T cells regulated by IL-6 signal is involved in superantigen-mediated CD4⁺ T cell death. *Int. Immunol.*, **21**, 73-80 (2008).
319. Kojima, M., Ashino, T., Yoshida, T., Iwakura, Y., Sekimoto, M., and Degawa, M. IL-1 regulates the *Cyp7a1* gene and serum total cholesterol level at steady state in mice. *Biochem. Biophys. Res. Commun.*, **379**, 239-242 (2008).
320. Chida, D., Sato, T., Sato, Y., Kubo, M., Yoda, T., Suzuki, H., and Iwakura, Y. Characterization of mice deficient in Melanocortin 2 receptor on a B6/Balbc mix background. *Mol. Cell. Endocrinol.*, **300**, 32-36 (2009).
321. Bai, Z. B., Hayasaka, H., Kobayashi, M., Li, W., Guo, Z., Jang, M.-H., Kondo, A., Choi, B., Iwakura, Y., and Miyasaka, M.

- CXCL12 promotes CCR7-dependent naïve T-cell trafficking to lymph nodes and Peyer's patches. *J. Immunol.*, **182**, 1287-1295 (2009).
322. Ishigame, H., Kakuta, S., Nagai, T., Nambu, A., Komiyama, Y., Kadoki, M., Tanahashi, Y., Akitsu, A., Kotaki, H., Sudo, K., Nakae, S., Sasakawa, C., and Iwakura, Y. Differential roles of IL-17A and IL-17F in host defense against mucoepithelial bacterial infection and allergic responses. *Immunity*, **30**, 108-119 (2009).
323. Yoneda, M., Suzuki, T., Nakamura, T., Ajima, R., Yoshida, Y., Kakuta, S., Sudo, K., Iwakura, Y., Shibutani, M., Mitsumori, K., Yokota, J., and Yamamoto, T. Deficiency of antiproliferative family protein Ana correlates with development of lung adenocarcinoma. *Cancer Sci.*, **100**, 225-232 (2009).
324. Lopez Kostka, S., Dinges, S., Griewank, K., Iwakura, Y., Udey, M. C., and Von Stebut, E. IL-17 promotes progression of cutaneous leishmaniasis in susceptible mice. *J. Immunol.*, **182**, 3039-3046 (2009).
325. Yamada, J., Hamuro, J., Fukushima, A., Ohteki, T., Terai, K., Iwakura, Y., Yagita, H., and Kinoshita, S. MHC-matched corneal allograft rejection in an IFN- γ /IL-17-independent manner in C57BL/6 mice. *Invest. Ophthalmol. Vis. Sci.*, **50**, 2139-2146 (2009).
326. Kuroishi, T., Kinbara, M., Sato, N., Tanaka, Y., Nagai, Y., Iwakura, Y., Endo, Y., and Sugawara, S. Biotin status affects nickel allergy via regulation of IL-1 β production in mice. *J. Nutrition*, **139**, 1031-1036 (2009).
327. Inoue, A., Setoguchi, K., Matsubara, Y., Okada, K., Sato, N., Iwakura, Y., Higuchi, O., and Yamanashi, Y. Dok-7 activates the muscle receptor kinase MuSK and shapes synapse formation. *Science Signaling* **2**, ra7 (2009).
328. Elkabets, M., Krelin, Y., Dotan, S., Cerwenka, A., Porgador, A., Lichtenstein, R. G., White, M. R., Zoller, M., Iwakura, Y., Dinarello, C. A., Voronov, E., and Apte, R. N. Host-derived Interleukin-1 α is

- important in determining the immunogenicity of 3-Methylcholantrene-tumor cells. *J. Immunol.*, **182**, 4874-4881 (2009).
329. Irie, N., Takada, Y., Watanabe, Y., Matsuzaki, Y., Naruse, C., Asano, M., Iwakura, Y., Suda, T., and Matsuo, K. Bidirectional signaling through EphrinA2-EphA2 enhances osteoclastogenesis and suppresses osteoblastogenesis. *J. Biol. Chem.*, **284**, 14637-14644 (2009).
 330. Di Paolo, N. C., Miao, E. A., Iwakura, Y., Kaja, M-K., Aderem, A., Flavell, R. A., Papayannopoulou, T., and Shayakhmetov, D. M. Virus sensing at the plasma membrane triggers interleukin-1 α -mediated pro-inflammatory macrophage response. *Immunity*, **31**, 110-121 (2009).
 331. Miyazato, A., Nakamura, K., Yamamoto, N., Mora-Montes, H. M., Tanaka, M., Abe, Y., Tanno, D., Inden, K., Gang, X., Ishii, K., Takeda, K., Akira, S., Saijo, S., Iwakura, Y., Adachi, Y., Ohno, N., Mitsutake, K., Gow, N. A., Kaku, M., and Kawakami, K. Toll-like receptor 9-dependent activation of myeloid dendritic cells by deoxynucleic acids from *Candida albicans*. *Infect. Immun.*, **77**, 3056-3064 (2009).
 332. Ohtaki, Y., Yamaguchi, K., Yu, Z., Kumamoto, H., Shimauchi, H., Iwakura, Y., Sugawara, S., and Endo, Y. Hepatic platelet accumulation in Fas-mediated hepatitis in mice. *Int. Immunopharmacol.*, **9**, 1071-1078 (2009).
 333. O'Connor, W. Jr., Kamanaka, M., Booth, C. J., Town, T., Nakae, S., Iwakura, Y., Kolls, J. K., and Flavell, R. A. A protective function for interleukin 17A in T cell-mediated intestinal inflammation. *Nat. Immunol.*, **10**, 603-609 (2009).
 334. Oseko, F., Yamamoto, T., Akamatsu, Y., Kanamura, N., Iwakura, Y., Imanishi, J., and Kita, M. IL-17 is involved in bone resorption in mouse periapical lesions. *Microbiol Immunol.*, **53**, 287-294 (2009).

335. Honore, P., Donnelly-Roberts, D., Namovic, M., Zhong, C., Wade, C., Chandran, P., Zhu, C., Carroll, W., Perez-Medrano, A., Iwakura, Y., and Jarvis, M. F. The antihyperalgesic activity of selective P2X7 receptor antagonist, A-839977, is lost in IL-1 α knockout mice. *Behav. Brain Res.*, **204**, 77-81 (2009).
336. Yi, T., Chen, Y., Wang, L., Du, G., Huang, D., Zhao, D., Johnstone, H., Young, J., Todorov, I., Umetsu, D. T., Chen, L., Iwakura, Y., Kandeel, F., Forman, S., and Zeng, D. Reciprocal differentiation and tissue-specific pathogenesis of Th1, Th2, and Th17 cells in graft versus host disease. *Blood*, **114**, 3101-3112 (2009).
337. Lindauer, M. L., Wong, J., Iwakura, Y., and Magun, B. E. Pulmonary inflammation triggered by Ricin toxin requires macrophages and IL-1 signaling. *J. Immunol.*, **183**, 1419-1426 (2009).
338. Shichita, T., Sugiyama, Y., Ooboshi, H., Sugimori, H., Nakagawa, R., Takada, I., Iwaki, T., Okada, Y., Iida, M., Cua, D. J., Iwakura, Y., and Yoshimura, A. Pivotal role of cerebral interleukin-17-producing $\gamma\delta$ T cells in the delayed phase of ischemic brain injury. *Nat. Med.*, **15**, 946-950 (2009).
339. Okiyama, N., Sugihara, T., Iwakura, Y., Yokozeki, H., Miyasaka, N., and Kohsaka, H. Therapeutic effects of interleukin-6 blockade in a murine model of polymyositis that does not require interleukin-17A. *Arth. Rheum.*, **60**, 2505-2512 (2009).
340. Horino, T., Matsumoto, T., Ishikawa, H., Kimura, S., Uramatsu, M., Tanabe, M., Tateda, K., Miyazaki, S., Aramaki, Y., Iwakura, Y., Yoshida, M., Onodera, S., and Yamaguchi, K. Interleukin-1 deficiency in combination with macrophage depletion increases susceptibility to *Pseudomonas aeruginosa* bacteremia. *Microbiol. Immunol.*, **53**, 502-511 (2009).
341. Taniguchi, S., Nakazawa, T., Tanimura, A., Kiyama, Y., Tezuka, T., Watabe, A. M., Katayama, N., Yokoyama, K., Inoue, T., Izumi-Nakaseko, H., Kakuta, S., Sudo, K., Iwakura, Y., Umemori, H.,

- Inoue, T., Murphy, N. P., Hashimoto, K., Kano, M., Manabe, T., and Yamamoto, T. Involvement of NMDAR2A tyrosine phosphorylation in depression-related behavior. *EMBO J.*, **28**, 3717-3729 (2009).
342. Horie, I., Abiru, N., Nagayama, Y., Kuriya, G., Saitoh, O., Ichikawa, T., Iwakura, Y., and Eguchi, K. T helper type 17 immune response plays an indispensable role for development of iodine-induced autoimmune thyroiditis in nonobese diabetic-H2h4 mice. *Endocrinology*, **150**, 5135-5142 (2009).
343. Lin, Y., Ritchie, S., Logar, A., Slight, S., Messmer, M., Rangel-Moreno, J., Gugliani, L., Alcorn, J. F., Strawbridge, H., Park, S. M., Onishi, R., Nyugen, N., Walter, M. J., Pociask, D., Randall, T. D., Gaffen, S. L., Iwakura, Y., Kolls, J. K., and Khader, S. A. Interleukin-17 is required for T helper 1 cell immunity and host resistance to the intracellular pathogen *Francisella tularensis*. *Immunity*, **31**, 799-810 (2009).
344. Tanaka, S., Yoshimoto, T., Naka, T., Nakae, S., Iwakura, Y., Cua, D., and Kubo, M. Natural occurring IL-17 producing T cells regulate the initial phase of neutrophil mediated airway responses. *J. Immunol.*, **183**, 7523-7530 (2009).
345. Faust, S. M., Lu, G., Marini, B. L., Zou, W., Gordon, D., Iwakura, Y., Laouar, Y., and Bishop, D. K. Role of T cell TGF β signaling and IL-17 in allograft acceptance and fibrosis associated with chronic rejection. *J. Immunol.*, **183**, 7297-7306 (2009).
346. Oyamada, A., Ikebata, H., Itsumi, M., Saiwai, H., Okada, S., Shimoda, K., Iwakura, Y., Nakayama, K. I., Iwamoto, Y., Yoshikai, Y., and Yamada, H. Tyrosine kinase 2 plays critical roles in the pathogenic CD4 T cell responses for the development of experimental autoimmune encephalomyelitis. *J. Immunol.*, **183**, 7539-7546 (2009).
347. Munoz, M., Heimesaat, M. M., Danker, K., Struck, D., Lohmann, U., Plickert, R., Bereswill, S., Fischer, A., Dunay, I. R., Wolk, K.,

- Loddenkemper, C., Krell, H.-W., Libert, C., Lund, L. R., Frey, O., Holscher, C., Iwakura, Y., Ghilardi, N., Ouyang, W., Kamradt, T., Sabat, R., and Liesenfeld, O. Interleukin (IL)-23 mediates *Toxoplasma gondii*-induced immunopathology in the gut via matrix metalloproteinase-2 and IL-22 but independent of IL-17. *J. Exp. Med.*, **206**, 3047-3059 (2009).
348. Doodes, P. D., Cao, Y., Hamel, K. M., Wang, Y., Rodeghero, R. I., Mikecz, K., Glant T. T., Iwakura, Y., and Finnegan, A. IFN- γ regulates the requirement for IL-17 in proteoglycan-induced arthritis. *J. Immunol.*, **184**, 1552-1559 (2010).
349. Madhur, M. S., Lob, H. E., McCann, L. A., Iwakura, Y., Blinder, Y., Guzik, T. J., and Harrison, D. G. Interleukin 17 promotes angiotensin II-induced hypertension and vascular dysfunction. *Hypertension*, **55**, 500-507 (2010).
350. Ohtaki, H., Yufu, S., Nakamachi, T., Satoh, K., Shimizu, A., Mori, H., Sato, A., Iwakura, Y., Matsunaga, M., and Shioda, S. Nucleoprotein diet ameliorates arthritis symptoms in mice transgenic for human T-cell leukemia virus type I (HTLV-I). *J. Clin. Biochem. Nutr.*, **46**, 93-104 (2010).
351. Kawa, K., Tsutsui, H., Uchiyama, R., Kato, J., Matsui, K., Iwakura, Y., Matsumoto, T., and Nakanishi, K. IFN- γ is a master regulator of endotoxin shock syndrome in mice primed with heat-killed *Propionibacterium acnes*. *Int. Immunol.*, **22**, 157-166 (2010).
352. Henry, T., Kirimanjeswara, G. S., Ruby, T., Jones, J. W., Peng, K., Perret, M., Ho, L., Sauer, J.-D., Iwakura, Y., Metzger, D. W., and Monack, D. M. Type I interferon signaling constrains IL-17A/F secretion by $\gamma\delta$ T cells during bacterial infection. *J. Immunol.*, **184**, 3755-3767 (2010).
353. Okae, H., and Iwakura, Y. Neural tube defects and impaired neural progenitor cell proliferation in *G β 1*-deficient mice. *Developmental Dynamics*, **239**, 1089-1101 (2010).

354. Okamoto Yoshida, Y., Umemura, M., Yahagi, A., O'Brien, R. L., Ikuta, K., Kishihara, K., Hara, H., Nakae, S., Iwakura, Y., and Matsuzaki, G. Essential role of IL-17A in the formation of a mycobacterial infection-induced granuloma in the lung. *J. Immunol.*, **184**, 4414-4422 (2010).
355. Kono, H., Karmarkar, D., Iwakura, Y., and Rock, K. L. Identification of the cellular sensor that stimulates the inflammatory response to sterile cell death. *J. Immunol.*, **184**, 4470-4478 (2010).
356. Isoda, K., Matsuki, T., Kondo, H., Iwakura, Y., and Ohsuzu, F. Deficiency of interleukin-1 receptor antagonist induces aortic valve disease in BALB/c mice. *Arterioscler. Thromb. Vasc. Biol.*, **30**, 708-715 (2010).
357. Itoh, S., Nakae, S., Axtell, R. C., Velotta, J. B., Kimura, N., Kajiwar, N., Iwakura, Y., Saito, H., Adachi, H., Steinman, L., Robbins, R. C., and Fishbein, M. P. IL-17 contributes to the development of chronic rejection in a murine-heart transplant model. *J. Clin. Immunol.*, **30**, 235-240 (2010).
358. Zhao, Y., Zhang, X., Guda, K., Lawrence, E., Sun, Q., Watanabe, T., Iwakura, Y., Asano, M., Wei, L., Yang, Z., Zheng, W., Dawson, D., Willis, J., Markowitz, S. D., Satake, M., and Wang, Z. Identification and functional characterization of paxillin as a target of protein tyrosine phosphatase receptor T. *Proc. Natl. Acad. Sci. USA*, **107**, 2592-2597 (2010).
359. St. Pierre, C. A., Chan, M., Iwakura, Y., Ayers, D. C., Kurt-Jones, E. A., and Finberg, R. W. Periprosthetic Osteolysis: Characterizing the innate immune response to titanium wear particles. *J. Orthopedic Res.*, **28**, 1418-1424 (2010).
360. Adamopoulos, I. E., Chao, C. C., Geissler, R., Lafage, D., Blumenschein, W., Iwakura, Y., McClanahan, T., and Bowman, E. P. Interleukin-17A upregulates receptor activator of NF- κ B on osteoclast precursors. *Arthritis Res. Ther.*, **12**, R29 (2010).

361. Kadoki, M., Choi, B. I., and Iwakura, Y. The mechanism of lipopolysaccharide-induced human immunodeficiency virus type I activation in transgenic mouse macrophages. *Int. Immunol.*, **22**, 469-478 (2010).
362. Kyuwa, S., Takagaki, S., Matsuyama, S., Taguchi, F., Saegusa, J., Iwakura, Y., Tagawa, Y., and Yoshikawa, Y. Characterization of a variant virus from ascitic fluid of subacute granulomatous serositis in interferon- γ -deficient C57BL/6 mice persistently infected with murine coronavirus strain JHM. *Viral Immunol.*, **23**, 437-442 (2010).
363. Teng, M. W. L., Andrews, D. M., McLaughlin, N., von Scheidt, B., Ngiow, S. F., Möller, A., Hill, G. R., Iwakura, Y., Oft, M., and Smyth, M. J. IL-23 suppresses innate immune response independently of IL-17A during carcinogenesis and metastasis. *Proc. Natl. Acad. Sci. USA*, **107**, 8328-8333 (2010).
364. Baldeviano, G. C., Barin, J. G., Talor, M. V., Srinivasan, S., Bedja, D., Zheng, D., Gabrielson, K., Iwakura, Y., Rose, N. R., and Cihakova, D. IL-17A is dispensable for myocarditis but essential for the progression to dilated cardiomyopathy. *Circulation Res.*, **106**, 1645-1655 (2010).
365. Saijo, S., Ikeda, S., Yamabe, K., Kakuta, S., Ishigame, H., Akitsu, A., Fujikado, N., Kusaka, T., Kubo, S., Chung, S., Komatsu, R., Miura, N., Adachi, Y., Ohno, N., Shibuya, K., Yamamoto, N., Kawakami, K., Yamasaki, S., Saito, T., Akira, S., and Iwakura, Y. Dectin-2 recognition of α -mannans and induction of Th17 differentiation is essential for host defense against *Candida albicans*. *Immunity*, **32**, 681-691 (2010).
366. Hill, G. R., Olver, S. D., Kuns, R. D., Varelias, A., Raffelt, N. C., Don, A. L. J., Markey, K. A., Wilson, Y. A., Smyth, M. J., Iwakura, Y., Tocker, J., Clouston, A., and MacDonald, K. P. A. Stem cell mobilization with G-CSF induces type-17 differentiation and promotes scleroderma. *Blood*, **116**, 819-828 (2010).

367. Gan, P. Y., Steinmetz, O. M., Tan, D. S., O'Sullivan, K. M., Ooi, J. D., Iwakura, Y., Kitching, A. R., and Holdsworth, S. R. Th17 cells promote autoimmune anti-myeloperoxidase glomerulonephritis. *J. Am. Soc. Nephrol.*, **21**, 925-931 (2010).
368. Cowley, S. C., Meierovics, A. I., Frelinger, J. A., Iwakura, Y., and Elkins, K. L. Lung CD4-CD8-double-negative T cells are prominent producers of IL-17A and IFN- γ during primary respiratory murine infection with *Francisella tularensis* live vaccine strain. *J. Immunol.*, **184**, 5791-5801 (2010).
369. Wakita, D., Sumida, K., Iwakura, Y., Nishikawa, H., Ohkuri, T., Chamoto, K., Kitamura, H., and Nishimura, T. Tumor-infiltrating IL-17-producing gammadelta T cells support the progression of tumor via promoting angiogenesis. *Eur. J. Immunol.*, **40**, 1927-1937 (2010).
370. Oh, I., Ozaki, K., Meguro, A., Hatanaka, K., Kadowaki, M., Matsu, H., Tatara, R., Sato, K., Iwakura, Y., Nakae, S., Sudo, K., Teshima, T., Leonard, W. J., and Ozawa, K. Altered effector CD4⁺ T cell function in IL-21R^{-/-} CD4⁺ T cell-mediated graft-versus-host-disease. *J. Immunol.*, **185**, 1920-1926 (2010).
371. Ashino, S., Wakita, D., Shiohama, Y., Iwakura, Y., Chamoto, K., Ohkuri, T., Kitamura, H., and Nishimura, T. A Th17-polarized cell population that has infiltrated the lung requires cells that convert to IFN- γ production in order to induce airway hyperresponsiveness. *Int. Immunol.*, **22**, 503-513 (2010).
372. Xiao, G., Miyazato, A., Abe, Y., Zhang, T., Nakamura, K., Inden, K., Tanaka, M., Tanno, D., Miyasaka, T., Ishii, K., Takeda, K., Akira, S., Saijo, S., Iwakura, Y., Adachi, Y., Ohno, N., Yamamoto, N., Kunishima, H., Hirakata, Y., Kaku, M., and Kawakami, K. Activation of myeloid dendritic cells by deoxynucleic acids from *Cordyceps sinensis* via a Toll-like receptor 9-dependent pathway. *Cellular Immunol.*, **263**, 241-250 (2010).

373. Miyazaki, Y., Hamano, S., Wang, S., Shimanoe, Y., Iwakura, Y., and Yoshida, H. IL-17 is necessary for host protection against acute-phase *Trypanosoma cruzi* infection. *J. Immunol.*, **185**, 1150-1157 (2010).
374. Nakajima, A., Matsuki, T., Komine, M., Asahina, A., Horai, R., Ishigame, H., Nakae, S., Kakuta, S., Saijo, S., and Iwakura, Y. TNF, but not IL-6 and IL-17, is crucial for the development of T cell-independent psoriasis-like dermatitis in *Il1rn*^{-/-} mice. *J. Immunol.*, **185**, 1887-1893 (2010).
375. Hadano, S., Yoshii, Y., Otomo, A., Kunita, R., Suzuki-Utsunomiya, K., Pan, L., Kakuta, S., Iwasaki, Y., Iwakura, Y., and Ikeda, J-E. Genetic background and gender effects on gross phenotypes in congenic lines of *ALS2/alsin*-deficient mice. *Neurosci. Res.*, **68**, 131-136 (2010).
376. Henningsson, L., Jirholt, P., Lindholm, C., Eneljung, T., Silverpil, E., Iwakura, Y., Linden, A., and Gjertsson, I. Interleukin-17A during local and systemic *Staphylococcus aureus*-induced arthritis in mice. *Infect. Immun.*, **78**, 3783-3790 (2010).
377. Dunne, A., Ross, P. J., Pospisilova, E., Masin, J., Meaney, A., Sutton, C. E., Iwakura, Y., Tschopp, J., Sebo, P., and Mills, K. H. Inflammasome activation by adnylate cyclase toxin directs Th17 responses and protection against *Bordetella pertussis*. *J. Immunol.*, **185**, 1711-1719 (2010).
378. Lee, Y.-M., Fujikado, N., Manaka, H., Yasuda, H., and Iwakura, Y. IL-1 plays an important role in the bone metabolism under physiological conditions. *Int. Immunol.*, **22**, 805-816 (2010).
379. Ikeda, U., Wakita, D., Ohkuri, T., Chamoto, K., Kitamura, H., Iwakura, Y., and Nishimura, T. 1 α , 25-Dihydroxyvitamin D(3) and all-trans retinoic acid synergistically inhibit the differentiation and expansion of Th17 cells. *Immunol. Lett.*, **134**, 7-16 (2010).
380. Horie, I., Abiru, N., Saitoh, O., Ichikawa, T., Iwakura, Y., Eguchi, K., and Nagayama, Y. Distinct role of T helper type 17 immune

response for Graves' hyperthyroidism in mice with different genetic background. *Autoimmunity*, **44**, 159-165 (2010).

381. Narita, K., Hu, D. L., Mori, F., Wakabayashi, K., Iwakura, Y., and Nakane, A. Role of IL-17A in cell-mediated protection against *Staphylococcus aureus* infection in mice immunized with fibrinogen binding domain of clumping factor A. *Infect. Immun.*, **78**, 4234-4242 (2010).
382. Chou, R. C., Kim, N. D., Sadik, C. D., Seung, E., Lan, Y., Byrne, M. H., Haribabu, B., Iwakura, Y., and Luster, A. D. Lipid-cytokine-chemokine cascade drives neutrophil recruitment in a murine model of inflammatory arthritis. *Immunity*, **33**, 266-278 (2010).
383. Vokaer, B., Van Rompaey, N., Lemaître, P. H., Lhommé, F., Kubjak, C., Benghiat, F. S., Iwakura, Y., Petein, M., Field, K. A., Goldman, M., Le Moine, A., and Charbonnier, L. M. Critical role of regulatory T cells in Th17-mediated minor antigen-disparate rejection. *J. Immunol.*, **185**, 3417-25 (2010).
384. Chung, S. H., Seki, K., Choi, B. I., Kimura, K. B., Ito, A., Fujikado, N., Saijo, S., and Iwakura, Y. CXC chemokine receptor 4 expressed in T cells plays an important role in the development of collagen-induced arthritis. *Arthritis Res. Ther.*, **12**, R188 (2010).
385. Lee, H. H., Meyer, E. H., Goya, S., Pichavant, M., Kim, H. Y., Bu, X., Umetsu, S. E., Jones, J. C., Savage, P. B., Iwakura, Y., Casasnovas, J. M., Kaplan, G., Freeman, G. J., Dekruyff, R. H., and Umetsu, D. T. Apoptotic cells activate NKT cells through T cell Ig-like mucin-like-1 resulting in airway hyperreactivity. *J. Immunol.*, **185**, 5225-5235 (2010).
386. Matsuwaki, T., Nishihara, M., Sato, T., Yoda, T., Iwakura, Y., and Chida, D. Functional hypothalamic amenorrhea due to increased CRH tone in melanocortin receptor2-deficient mice. *Endocrinology*, **151**, 5489-5496 (2010).

387. Ishida, H., Matsuzaki-Moriya, C., Imai, T., Yanagisawa, K., Nojima, Y., Suzue, K., Hirai, M., Iwakura, Y., Yoshimura, A., Hamano, S., Shimokawa, C., and Hisaeda, H. Development of experimental cerebral malaria is independent of IL-23 and IL-17. *Biochem. Biophys. Res. Commun.*, **402**, 790-795 (2010).
388. Sato, Y., Suzuki, H., Sato, T., Suda, T., Yoda, T., Iwakura, Y., and Chida, D. The role of endogenous glucocorticoid in lymphocyte development in melanocortin receptor 2-deficient mice. *Biochem. Biophys. Res. Commun.*, **403**, 253-257 (2010).
389. Sun, X., Yamada, H., Shibata, K., Muta, H., Tani, K., Podack, E. R., Iwakura, Y., and Yoshikai, Y. CD30 ligand is a target for a novel biological therapy against colitis associated with Th17 responses. *J. Immunol.*, **185**, 7671-7680 (2010).