WELL-POSEDNESS FOR THE NONLINEAR SCHRÖDINGER EQUATION WITH DERIVATIVE NONLINEARITY

HIDEO TAKAOKA
MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY,
SENDAI, 980-8578, JAPAN

This note concerns the well-posedness for the Cauchy problem of derivative nonlinear Schrödinger equation:

$$i\partial_t u + \partial_x^2 u = i\lambda\partial_x(|u|^2 u), \quad (t,x) \in \mathbb{R}^2, \tag{1}$$

$$u(0,x) = u_0(x), \quad x \in \mathbb{R}, \tag{2}$$

where the unknown function u is a complex valued function of $(t,x) \in \mathbb{R}^2$ and $\lambda \in \mathbb{R}$. The derivative nonlinear Schrödinger equation is a model of the propagation of circularly polarized Alfvèn waves in magnetized plasma with a constant external magnetic field [15, 16].

Here we consider the well-posedness, which means the existence, the uniqueness, the persistency property ($u_0 \in X \Rightarrow u(t) \in X$) and the continuous dependence of solution on data ($u_0 \to v_0 \Rightarrow u(t) \to v(t)$). We say the global (resp. local) well-posedness, if the well-posedness holds for all time (resp. a positive time interval).

When we try to show the well-posedness by the contraction argument, we meet with the derivative loss stemming from the derivative nonlinearity of equation (1). In [9, 10, 11, 17], the H^1 global well-posedness for small initial data was shown, where the proof uses two gauge transformations. The strategy of argument in [9, 10, 11, 17] is to reduce the original equation (1) to equations without derivative nonlinearity.

The well-posedness below energy is considered in [18, 20]. In [18] the $H^{\frac{1}{2}}$ local well-posedness was proven. The proof of [18] was a suitable combination of the gauge transformation and the Fourier restriction norm method [1, 12]. The transformation [9, 10, 11, 17, 18]

$$v(t,x) = \exp\left(-i\lambda\int_{-\infty}^{x}|u(t,y)|^2 dy\right)u(t,x), \tag{3}$$

reduces (1) to the following:

$$i\partial_t v + \partial_x^2 v = -i\lambda v^2\partial_x\overline{v} - \frac{\lambda^2}{2}|v|^4 v. \tag{4}$$

Solutions of the Cauchy problem corresponding to (5) satisfy the conservation law:

$$E_1(v(t)) = \|\partial_x v(t)\|_{L^2_x}^2 - \frac{\lambda}{2}\Im\langle |v(t)|^2 v(t), \partial_x v(t)\rangle_{L^2_x}. \tag{5}$$

As compared to the equation (1), $|v|^4 v$ appears and $v^2\partial_x\overline{v}$ remains in (3), but the term of derivative type $|v|^2\partial_x v$ in (1) is removed here. The term $|v|^2\partial_x v$ is difficult to handle, while the proof in [18] can control the terms $v^2\partial_x\overline{v}$ and $|v|^4 v$.

Anyway, it is natural to try to extend the above local solution below energy to global one. But it is not obvious, because it seems difficult to give the a priori bound of solution by the usual use of conservation laws (cf. (5), L^2-conservation, etc ...). The above problem is investigated in [20]. The proof of [20] is based on the argument in [3]. In [3] Bourgain shows global existence results for the 2D-NLS in weaker spaces than in the energy space (see also [4, 5, 3D-NLS], [6, 7, KdV], [8, mKdV], [13, 14, NLW], [19, 23, KP-2]). His strategy is a general scheme as follows. Let V_t and $V(t)$ denote the flow maps of nonlinear and linear equations, that is, $V_t u_0$ and $V(t)u_0$ are solutions of nonlinear and linear equations, respectively. Let X and Y be Banach spaces such that $X \subsetneq Y$, where X and Y correspond to conservation law spaces and the initial data space, respectively. His strategy is that if $(V_t - V(t))u_0 \in X$ for all time t, we have the global well-posedness in Y. Roughly speaking, his argument aims to estimate the high Sobolev norm of solution by the low Sobolev norm, which controls the transportation of energy from the low frequency region to the high frequency region. He applied this argument to NLS with $X = H^1$ and $Y = H^s$ for some $0 < s < 1$ in [3]. However in equations of derivative type, the derivative loss stems from the derivative nonlinearity. Moreover we have to regain more than one derivatives for the usage of Bourgain's argument. Here we combine the above argument of Bourgain with the Fourier restriction norm method observed in [18].

This talk devotes to the above results of [18, 20]. More precisely, we are to exhibit the following two theorems.

Theorem 1 ([18]). *Let $s \geq \frac{1}{2}$. Then the Cauchy problem (1)-(2) is locally well-posed in H^s.*

Theorem 2 ([20]). *The Cauchy problem (1)-(2) is globally well-posed in H^s for $s_0 < s < 1$ for appropriate $s_0 \in (\frac{1}{2}, 1)$, provided $\|u_0\|_{L^2} < \sqrt{\frac{2\pi}{|\lambda|}}$.*

Concerning the number s_0 in Theorem 2, we shall observe in the talk.

References

[1] SchrJ. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, part I: Schrödinger equation, part II: The KdV equation*, Geom. Funct. Anal. 3 (1993), 107–156, 209–262.

[2] J. Bourgain, *Periodic Korteweg de Vries equation with measure as initial data*, Sel. Math. New Ser. 3 (1997), 115–159.

[3] J. Bourgain, *Refinement of Strichartz inequality and applications to 2D-NLS with critical nonlinearity*, Internat. Math. Res. Notices 5 (1998), 253–283.

[4] J. Bourgain, *Global wellposedness of defocusing critical nonlinear Schrödinger equation in the radial case*, J. Amer. Math. Soc. 12 (1999), 145–171.

[5] J. Bourgain, *Scattering in the energy space and below for 3D NLS*, J. Analyse Math. 75 (1998), 267–297.

[6] J. Colliander, G. Staffilani and H. Takaoka, *Global well-posedness for KdV below* L^2, Math. Res. Lett. 6 (1999), 775–798.

[7] J. Colliander, G. Staffilani, M. Keel, H. Takaoka and T. Tao, *Global well-posedness of the KdV equation below the* L^2 *norm*, in preparation

[8] G. Fonseca, F. Linares and G. Ponce, *Global well-posedness for the modified Korteweg-de Vries equation*, Comm. partial Differential Equations 24 (1999), 683–705.

[9] N. Hayashi, *The initial value problem for the derivative nonlinear ödinger equation in the energy space*, Nonlinear Anal. 20 (1993), 823–833.

[10] N. Hayashi and T. Ozawa, *On the derivative nonlinear Schrödinger equation*, Phys. D 55 (1992), 14–36.

[11] N. Hayashi and T. Ozawa, *Finite energy solutions of nonlinear Schrödinger equations of derivative type*, SIAM J. Math. Anal. 25 (1994), 1488–1503.

[12] C. E. Kenig, G. Ponce and L. Vega, *A bilinear estimate with applications to the KdV equation*, J. Amer. Math. Soc. 9 (1996), 573–603.

[13] C. E. Kenig, G. Ponce and L. Vega, *Global well-posedness for semi-linear wave equations*, preprint.

[14] M. Keel and T. Tao, *Local and global well-posedness of wave maps of* $\mathbb{R}^{1+1}$ *for rough data*, Internat. Math. Res. Notices 21 (1998), 1117–1156.

[15] W. Mio, T. Ogino, K. Minami and S. Takeda, *Modified nonlinear Schrödinger equation for Alfvèn waves propagating along the magnetic field in cold plasma*, J. Phys. Soc. Japan 41 (1976), 265–271.

[16] E. Mjϕlhus, *On the modulational instability of hydromagnetic waves parallel to the magnetic field*, J. Plasma Phys. 16 (1976), 321–334.

[17] T. Ozawa, *On the nonlinear Schödinger equations of derivative type*, Indiana Univ. Math. J. 45 (1996), 137–163.

[18] H. Takaoka, *Well-posedness for the one dimensional nonlinear Schrödinger equation with the derivative nonlinearity*, Adv. Differential Equations 4 (1999), 561–580.

[19] H. Takaoka, *Global well-posedness for the Kadomtsev-Petviashvili II equation*, Discrete Contin. Dynam. Systems 6 (2000), 483–499.

[20] H. Takaoka, *Global well-posedness for the Schrödinger equations with derivative nonlinearity below energy*, preprint.

[21] M. Tsutsumi and I. Fukuda, *On solutions of the derivative nonlinear Schrödinger equation: existence and uniqueness theorem*, Funkcial. Ekvac. 23 (1980), 259–277.

[22] M. Tsutsumi and I. Fukuda, *On solutions of the derivative nonlinear Schrödinger equation II*, Funkcial. Ecvac. 24 (1981), 85–94.

[23] N. Tzvetkov, *Global low regularity solutions for Kadomtsev-Petviashvili equation*, Diff. Int. Eq. (to appear).