

# Oscillatory bifurcation for semilinear ordinary differential equations

Tetsutaro Shibata

Laboratory of Mathematics, Graduate School of Engineering,

Hiroshima University, Higashi-Hiroshima, 739-8527, Japan

tshibata@hiroshima-u.ac.jp

We consider the bifurcation problem

$$\begin{aligned} -u''(t) &= \lambda(u(t) + g(u(t))), & x \in I := (-1, 1), \\ u(t) &> 0, & t \in I, \\ u(-1) &= u(1) = 0. \end{aligned}$$

Here,  $\lambda > 0$  is a bifurcation parameter. The typical example of  $g(u)$  is  $g_1(u) := \sin \sqrt{u}$ . It is well known that under the suitable conditions on  $g(u)$ ,  $\lambda$  is parameterized by the maximum norm  $\alpha = \|u_\lambda\|_\infty$  of the solution  $u_\lambda$  corresponding to  $\lambda$  and is written as  $\lambda = \lambda(g, \alpha)$ . It should be mentioned that if  $g(u) = g_1(u) = \sin \sqrt{u}$ , then this problem has been proposed in Cheng (2002) as an example which has arbitrary many solutions near the line  $\lambda = \pi^2/4$ . In this talk, we first show that the bifurcation diagram of  $\lambda(g_1, \alpha)$  intersects the line  $\lambda = \pi^2/4$  infinitely many times by establishing the precise asymptotic formulas for  $\lambda(g_1, \alpha)$  as  $\alpha \rightarrow \infty$ . Secondly, we generalize the results above and treat the other cases, which produce the bifurcation curves intersecting the line  $\lambda = \pi^2/4$  infinitely many times.