

On supermagic coverings of fans and ladders

A.A.G. Ngurah, A.N.M. Salman and I W. Sudarsana

(Received September 22, 2009; Revised May 16, 2010)

Abstract. A simple graph G admits an H -covering if every edge in $E(G)$ belongs to a subgraph of G isomorphic to H . The graph G is said to be H -magic if there exists a bijection $f : V(G) \cup E(G) \rightarrow \{1, 2, 3, \dots, |V(G) \cup E(G)|\}$ such that for every subgraph H' of G isomorphic to H , $\sum_{v \in V(H')} f(v) + \sum_{e \in E(H')} f(e)$ is constant. Additionally, G is said to be H -supermagic if $f(V(G)) = \{1, 2, 3, \dots, |V(G)|\}$. In this paper, we study H -supermagic labelings of two classes of connected graph namely fans and ladders.

AMS 2000 *Mathematics Subject Classification.* 05C78.

Key words and phrases. Covering, cycle, fan, H -supermagic labeling, ladder.

§1. Introduction

We consider finite, undirected and simple graphs. The vertex and edge sets of a graph G are denoted by $V(G)$ and $E(G)$, respectively. Let H be a graph. An *edge-covering* of G is a family of subgraphs H_1, \dots, H_k such that each edge of $E(G)$ belongs to at least one of the subgraphs H_i , $1 \leq i \leq k$. If every H_i is isomorphic to a given graph H then we say that G admits an H -covering. Suppose G admits an H -covering. A total labeling $f : V(G) \cup E(G) \rightarrow \{1, 2, 3, \dots, |V(G) \cup E(G)|\}$ is said an H -magic labeling of G if for every subgraph H' of G isomorphic to H , $\sum_{v \in V(H')} f(v) + \sum_{e \in E(H')} f(e)$ is constant. An H -magic labeling f is said an H -supermagic labeling if $f(V(G)) = \{1, 2, 3, \dots, |V(G)|\}$. A graph that admits H -(super)magic labeling is called H -(super)magic. The sum of all vertex labels and all edge labels on H (under a labeling f) is denoted by $\sum f(H)$. In Figure 1, we show C_4 -magic and C_4 -supermagic labelings of L_4 .

The H -supermagic labeling was first introduced by Gutiérrez and Lladó [5] in 2005. They considered star-supermagic and path-supermagic labelings of some graphs. In [8], Lladó and Moragas gave C_n -supermagic labelings of




Figure 1: (a). a C_4 -magic labeling of L_4 (b). a C_4 -supermagic labeling of L_4

wheels, windmills, prisms and theta graphs. Cycles-supermagic labeling of chain graphs kC_n -snake, triangle ladders TL_n , grids $P_m \times P_n$, for $n = 2, 3, 4, 5$, and books B_n can be found in [13]. Maryati et al. [9] proved that some classes of trees such as subdivision of stars, shrubs, and banana tree graphs are P_h -supermagic for some h and prove that certain shackles and amalgamations of a connected graph H are H -supermagic [10].

For $H \cong P_2$, an H -supermagic graph is also called a *super edge-magic graph*. The notion of a super edge-magic graph was introduced by Enomoto et al. [2] as a particular type of edge-magic graph given by Kotzig and Rosa [6]. There are many graphs that have been proved to be (super) edge-magic graphs, see for instance [3, 11, 12, 14, 15]. For further information about (super) edge-magic graphs, see [4]. The H -magic labeling is related to a face-magic labeling of a plane graph introduced by Lih [7]. A total labeling f of a plane graph is said to be *face-magic* if for every positive integer s , all s -sided faces have the same weight. The weight of a face under a labeling f is the sum of labels carried by the edges and vertices surrounding it. Lih [7] allows different weights for different s . If a plane graph G contains only n -sided faces, then face-magic labeling of G is also C_n -magic labeling. Other results about this labeling can be found in [1].

In this paper, we study C_m and F_m -supermagic labelings of fans F_n , and C_m and L_m -supermagic labelings of ladders L_n for all possible values of m and n .

§2. Supermagic coverings of fans

In this section we consider C_m and F_m -supermagic labelings of the fans F_n . We define the fans $F_n \cong P_n + \{c\}$ as a graph with

$$V(F_n) = \{c, x_i | i = 1, 2, 3, \dots, n\}$$

and

$$E(F_n) = \{cx_i | i = 1, 2, 3, \dots, n\} \cup \{x_i x_{i+1} | i = 1, 2, 3, \dots, n-1\}.$$

In [7], Lih proved that F_n is C_3 -supermagic for every n except $n \equiv 2 \pmod{4}$. Furthermore, Ngurah et al. [13] proved that F_n is C_3 -supermagic for any $n \geq 2$. In the following theorem, we show that F_n is C_m -supermagic for any integer $4 \leq m \leq \lfloor \frac{n+4}{2} \rfloor$.

Theorem 1. *Let $n \geq 4$ be a positive integer. Then the fan F_n is C_m -supermagic for any integer $4 \leq m \leq \lfloor \frac{n+4}{2} \rfloor$.*

Proof. First, label every vertex in the following way.

- Label the vertex c with 1.

Case 1: $n \equiv 0 \pmod{m-1}$

- Label $x_1, x_m, x_{2m-1}, x_{3m-2}, x_{4m-3}, \dots, x_{n-m+2}$ with $2, 3, 4, 5, 6, \dots, \frac{n}{m-1} + 1$, respectively.
- For $1 \leq k \leq m-2$, label $x_{1+k}, x_{m+k}, x_{2m+k-1}, x_{3m+k-2}, x_{4m+k-3}, \dots, x_{n-m+k+2}$ with $k(\frac{n}{m-1}) + 2, k(\frac{n}{m-1}) + 3, k(\frac{n}{m-1}) + 4, k(\frac{n}{m-1}) + 5, k(\frac{n}{m-1}) + 6, \dots, (k+1)(\frac{n}{m-1}) + 1$, respectively.

Case 2: $n \equiv 1 \pmod{m-1}$

- Label $x_1, x_m, x_{2m-1}, x_{3m-2}, x_{4m-3}, \dots, x_{n-m+1}, x_n$ with $2, 3, 4, 5, 6, \dots, \frac{n-1}{m-1} + 1, \frac{n-1}{m-1} + 2$, respectively.
- For $1 \leq k \leq m-2$, label $x_{1+k}, x_{m+k}, x_{2m+k-1}, x_{3m+k-2}, x_{4m+k-3}, \dots, x_{n-m+k+1}$ with $k(\frac{n-1}{m-1}) + 3, k(\frac{n-1}{m-1}) + 4, k(\frac{n-1}{m-1}) + 5, k(\frac{n-1}{m-1}) + 6, k(\frac{n-1}{m-1}) + 7, \dots, (k+1)(\frac{n-1}{m-1}) + 2$, respectively.

Case 3: $n \equiv t \pmod{m-1}$, where $t = 2, 3, 4, \dots, m-2$

- Label $x_1, x_m, x_{2m-1}, x_{3m-2}, x_{4m-3}, \dots, x_{n-m-t+2}, x_{n-t+1}$ with $2, 3, 4, 5, 6, \dots, \frac{n-t}{m-1} + 1, \frac{n-t}{m-1} + 2$, respectively.
- Label $x_{n-t+2}, x_{n-t+3}, x_{n-t+4}, \dots, x_n$ with $2(\frac{n-t}{m-1}) + 3, 3(\frac{n-t}{m-1}) + 4, 4(\frac{n-t}{m-1}) + 5, \dots, t(\frac{n-t}{m-1}) + t + 1$, respectively.
- For $1 \leq k \leq t$, label $x_{1+k}, x_{m+k}, x_{2m+k-1}, x_{3m+k-2}, x_{4m+k-3}, \dots, x_{n-m-t+k+2}$ with $\gamma_1^k + 2, \gamma_1^k + 3, \gamma_1^k + 4, \gamma_1^k + 5, \gamma_1^k + 6, \dots, \gamma_1^k + (\frac{n-t}{m-1} + 1)$, respectively, where $\gamma_1^k = k(\frac{n-t}{m-1} + 1)$.

- For $t+1 \leq k \leq m-2$, label $x_{1+k}, x_{m+k}, x_{2m+k-1}, x_{3m+k-2}, x_{4m+k-3}, \dots, x_{n-m-t+k+2}$ with $\gamma_2^k + 2, \gamma_2^k + 3, \gamma_2^k + 4, \gamma_2^k + 5, \gamma_2^k + 6, \dots, \gamma_2^k + (\frac{n-t}{m-1}) + 1$, respectively, where $\gamma_2^k = k(\frac{n-t}{m-1}) + t$.

Next, label every edge as follows.

- For $1 \leq i \leq n$, label cx_i with $3n + 1 - i$.

For labeling the remaining edges, let $e_i = x_i x_{i+1}$, $1 \leq i \leq n-1$, and let $q = n-1$.

Case 1: $q \equiv 0 \pmod{m-2}$

- Label $e_1, e_{m-1}, e_{2m-3}, e_{3m-5}, e_{4m-7}, \dots, e_{q-m+3}$ with $n+2, n+3, n+4, n+5, n+6, \dots, n + \frac{q}{m-2} + 1$, respectively.
- For $1 \leq k \leq m-3$, label $e_{1+k}, e_{m-1+k}, e_{2m-3+k}, e_{3m-5+k}, e_{4m-7+k}, \dots, e_{q-m+3+k}$ with $\gamma_3^k + 2, \gamma_3^k + 3, \gamma_3^k + 4, \gamma_3^k + 5, \gamma_3^k + 6, \dots, \gamma_3^k + (\frac{q}{m-2}) + 1$, respectively, where $\gamma_3^k = k(\frac{q}{m-2}) + n$.

Case 2: $q \equiv 1 \pmod{m-2}$

- Label $e_1, e_{m-1}, e_{2m-3}, e_{3m-5}, e_{4m-7}, \dots, e_q$ with $n+2, n+3, n+4, n+5, n+6, \dots, n + \frac{q-1}{m-2} + 2$, respectively.
- For $1 \leq k \leq m-3$, label $e_{1+k}, e_{m-1+k}, e_{2m-3+k}, e_{3m-5+k}, e_{4m-7+k}, \dots, e_{q-m+2+k}$ with $\gamma_4^k + 3, \gamma_4^k + 4, \gamma_4^k + 5, \gamma_4^k + 6, \gamma_4^k + 7, \dots, \gamma_4^k + (\frac{q-1}{m-2}) + 2$, respectively, where $\gamma_4^k = k(\frac{q-1}{m-2}) + n$.

Case 3: $q \equiv t \pmod{m-2}$, where $t = 2, 3, 4, \dots, m-3$

- Label $e_1, e_{m-1}, e_{2m-3}, e_{3m-5}, e_{4m-7}, \dots, e_{q-t+1}$ with $n+2, n+3, n+4, n+5, n+6, \dots, n + \frac{q-t}{m-2} + 2$, respectively.
- Label $e_{q-t+2}, e_{q-t+2}, e_{q-t+3}, \dots, e_q$ with $n+2(\frac{q-t}{m-2})+3, n+3(\frac{q-t}{m-2})+4, n+4(\frac{q-t}{m-2})+5, \dots, n+t(\frac{q-t}{m-2})+t+1$, respectively.
- For $1 \leq k \leq t$, label $e_{1+k}, e_{m-1+k}, e_{2m-3+k}, e_{3m-5+k}, e_{4m-7+k}, \dots, e_{q-t-m+3+k}$ with $\gamma_5^k + 2, \gamma_5^k + 3, \gamma_5^k + 4, \gamma_5^k + 5, \gamma_5^k + 6, \dots, \gamma_5^k + \frac{q-t}{m-2} + 1$, respectively, $\gamma_5^k = k(\frac{q-t}{m-2} + 1) + n$.
- For $t+1 \leq k \leq m-3$, label $e_{1+k}, e_{m-1+k}, e_{2m-3+k}, e_{3m-5+k}, e_{4m-7+k}, \dots, e_{q-t-m+3+k}$ with $\gamma_6^k + 2, \gamma_6^k + 3, \gamma_6^k + 4, \gamma_6^k + 5, \gamma_6^k + 6, \dots, \gamma_6^k + \frac{q-t}{m-2} + 1$, respectively, $\gamma_6^k = k(\frac{q-t}{m-2}) + n + t$.

Let us denote the total labeling defined above by h . It can be checked that $h(V(F_n)) = \{1, 2, 3, \dots, n+1\}$; for $1 \leq i \leq n-m+1$, $h(x_i) = h(x_{i+m-1}) - 1$, $h(x_i x_{i+1}) = h(x_{i+m-2} x_{i+m-1}) - 1$, and $h(cx_i) + h(cx_{i+m-2}) = h(cx_{i+1}) + h(cx_{i+m-1}) + 2$.

For $1 \leq i \leq n-m+2$, let $C_m^{(i)}$ be the subcycle of F_n with $V(C_m^{(i)}) = \{c, x_j | i \leq j \leq i+m-2\}$ and $E(C_m^{(i)}) = \{cx_i, cx_{i+m-2}\} \cup \{x_j x_{j+1} | i \leq j \leq i+m-3\}$. It is easy to verify that for $1 \leq i \leq n-m+1$, $\sum h(C_m^{(i)}) = \sum h(C_m^{(i+1)})$. Thus, for $1 \leq i \leq n-m+2$, $\sum h(C_m^{(i)})$ is constant. Hence, F_n is C_m -supermagic for any integer $4 \leq m \leq \lfloor \frac{n+4}{2} \rfloor$. \square

Next, we consider fan-supermagic labelings of fan. Notice that F_n is $C_3 \cong F_2$ -supermagic [13] and F_n is trivially F_n -supermagic. In the following theorem, we show that F_n is F_m -supermagic for all remaining possible values of m .

Theorem 2. *Let $n \geq 4$ be a positive integer. The fan F_n is F_m -supermagic for every integer $3 \leq m \leq n-1$.*

Proof. Define a total labeling of F_n as follows.

- For $1 \leq i \leq n-1$, label $x_i x_{i+1}$ with $n+1+i$.
- For $1 \leq i \leq n$, label cx_i with $3n+1-i$.
- Label the vertex c with 1.

For the remaining vertices, we consider three following cases.

Case 1: $n \equiv 0 \pmod{m}$

- Label $x_1, x_{m+1}, x_{2m+1}, x_{3m+1}, \dots, x_{n-m+1}$ with $2, 3, 4, 5, \dots, \frac{n}{m} + 1$, respectively.
- For $1 \leq k \leq m-1$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m+1+k}$ with $k(\frac{n}{m}) + 2, k(\frac{n}{m}) + 3, k(\frac{n}{m}) + 4, k(\frac{n}{m}) + 5, \dots, (k+1)(\frac{n}{m}) + 1$, respectively.

Case 2: $n \equiv 1 \pmod{m}$

- Label $x_1, x_{m+1}, x_{2m+1}, x_{3m+1}, \dots, x_{n-m}, x_n$ with $2, 3, 4, 5, \dots, \frac{n-1}{m} + 1, \frac{n-1}{m} + 2$, respectively.
- For $1 \leq k \leq m-1$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m+k}$ with $k(\frac{n-1}{m}) + 3, k(\frac{n-1}{m}) + 4, k(\frac{n-1}{m}) + 5, k(\frac{n-1}{m}) + 6, \dots, (k+1)(\frac{n-1}{m}) + 2$, respectively.

Case 3: $n \equiv t \pmod{m}$, where $t = 2, 3, 4, \dots, m-1$

- Label $x_1, x_{m+1}, x_{2m+1}, x_{3m+1}, \dots, x_{n-t+1}$ with $2, 3, 4, 5, \dots, \frac{n-t}{m} + 2$, respectively.
- Label $x_{n-t+2}, x_{n-t+3}, x_{n-t+4}, \dots, x_n$ with $2(\frac{n-t}{m}) + 3, 3(\frac{n-t}{m}) + 4, 4(\frac{n-t}{m}) + 5, \dots, t(\frac{n-t}{m}) + t + 1$, respectively.
- For $1 \leq k \leq t$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m-t+1+k}$ with $k(\frac{n-t}{m} + 1) + 2, k(\frac{n-t}{m} + 1) + 3, k(\frac{n-t}{m} + 1) + 4, k(\frac{n-t}{m} + 1) + 5, \dots, (k + 1)(\frac{n-t}{m} + 1)$, respectively.
- For $t + 1 \leq k \leq m - 1$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m-t+1+k}$ with $k(\frac{n-t}{m}) + t + 2, k(\frac{n-t}{m}) + t + 3, k(\frac{n-t}{m}) + t + 4, k(\frac{n-t}{m}) + t + 5, \dots, (k + 1)(\frac{n-t}{m}) + t + 1$, respectively.

Denote the total labeling defined above by f . It can be checked that $f(V(F_n)) = \{1, 2, 3, \dots, n + 1\}$; for $1 \leq i \leq n - m + 1$,

$$\sum_{j=i}^{i+m-1} f(x_j) = -1 + \sum_{j=i+1}^{i+m} f(x_j),$$

$$\sum_{j=i}^{i+m-2} f(x_j x_{j+1}) = 1 - m + \sum_{j=i+1}^{i+m-1} f(x_j x_{j+1}),$$

and

$$\sum_{j=i}^{i+m-1} f(cx_j) = m + \sum_{j=i+1}^{i+m} f(cx_j).$$

For $1 \leq i \leq n - m + 1$, let $F_m^{(i)}$ be the subfan of F_n with $V(F_m^{(i)}) = \{c, x_j | i \leq j \leq i + m - 1\}$ and $E(F_m^{(i)}) = \{x_j x_{j+1} | i \leq j \leq i + m - 2\} \cup \{cx_j | i \leq j \leq i + m - 1\}$. It is a routine procedure to verify that for $1 \leq i \leq n - m$, $\sum f(F_m^{(i)}) = \sum f(F_m^{(i+1)})$. So, f is an F_m -supermagic labeling of F_n . Hence, F_n is F_m -supermagic. \square

In Figure 2, we show a C_4 -supermagic labeling of F_8 and an F_4 -supermagic labeling of F_{10} as defined in the proof of Theorems 1 and 2, respectively.

§3. Supermagic coverings of ladders

Let $L_n \cong P_n \times P_2$ denote the ladder of order $2n$ and size $3n - 2$. Clearly L_n admits a cycle covering of some even order. As a direct consequence of Lladó and Moragas's result (see Theorem 7 [8]), L_n is C_4 -supermagic for odd n . Later, Ngurah et al. [13] solved for the remaining cases. In the next theorem, we show that L_n is also C_{2m} -supermagic for the remaining possible values of m .




Figure 2: (a). a C_4 -supermagic labeling of F_8 (b). an F_4 -supermagic labeling of F_{10}

Theorem 3. *Let $n \geq 4$ be a positive integer. Then the ladder L_n is C_{2m} -supermagic for every integer $3 \leq m \leq \lfloor \frac{n}{2} \rfloor + 1$.*

Proof. First, let L_n be a graph with

$$V(L_n) = \{x_i, y_i | 1 \leq i \leq n\}$$

and

$$E(L_n) = \{x_i x_{i+1}, y_i y_{i+1} | 1 \leq i \leq n-1\} \cup \{x_i y_i | 1 \leq i \leq n\}.$$

Next, label every edge in the following way.

- For $1 \leq i \leq n-1$, label $x_i x_{i+1}$ with $2n+i$.
- For $1 \leq i \leq n-1$, label $y_i y_{i+1}$ with $4n-1-i$.
- For $1 \leq i \leq n$, label $x_i y_i$ with $5n-1-i$.

Label every vertex in the following way.

Case 1: $n \equiv 0 \pmod{m}$

- Label $x_1, x_{m+1}, x_{2m+1}, x_{3m+1}, \dots, x_{n-m+1}$ with $1, 2, 3, 4, \dots, \frac{n}{m}$, respectively.
- For $1 \leq k \leq m-1$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m+1+k}$ with $k(\frac{n}{m})+1, k(\frac{n}{m})+2, k(\frac{n}{m})+3, k(\frac{n}{m})+4, \dots, (k+1)(\frac{n}{m})$, respectively.

Case 2: $n \equiv 1 \pmod{m}$

- Label $x_1, x_{m+1}, x_{2m+1}, x_{3m+1}, \dots, x_{n-m}, x_n$ with $1, 2, 3, 4, \dots, (\frac{n-1}{m}), (\frac{n-1}{m})+1$, respectively.
- For $1 \leq k \leq m-1$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m+k}$ with $k(\frac{n-1}{m})+2, k(\frac{n-1}{m})+3, k(\frac{n-1}{m})+4, k(\frac{n-1}{m})+5, \dots, (k+1)(\frac{n-1}{m})+1$, respectively.

Case 3: $n \equiv t \pmod{m}$, where $t = 2, 3, 4, \dots, m-1$

- Label $x_1, x_{m+1}, x_{2m+1}, x_{3m+1}, \dots, x_{n-t+1}$ with $1, 2, 3, 4, \dots, (\frac{n-t}{m})+1$, respectively.
- Label $x_{n-t+2}, x_{n-t+3}, \dots, x_n$ with $2(\frac{n-t}{m}+1), 3(\frac{n-t}{m}+1), \dots, t(\frac{n-t}{m}+1)$, respectively.
- For $1 \leq k \leq t$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m-t+1+k}$ with $k(\frac{n-t}{m}+1)+1, k(\frac{n-t}{m}+1)+2, k(\frac{n-t}{m}+1)+3, k(\frac{n-t}{m}+1)+4, \dots, k(\frac{n-t}{m}+1)+\frac{n-t}{m}$, respectively.
- For $t+1 \leq k \leq m-1$, label $x_{1+k}, x_{m+1+k}, x_{2m+1+k}, x_{3m+1+k}, \dots, x_{n-m-t+1+k}$ with $k(\frac{n-t}{m})+t+1, k(\frac{n-t}{m})+t+2, k(\frac{n-t}{m})+t+3, k(\frac{n-t}{m})+t+4, \dots, (k+1)(\frac{n-t}{m})+t$, respectively.

Finally, for $1 \leq i \leq n$, label y_i with $n +$ (the label of x_i).

Let us denote the total labeling defined above by f . It can be checked that $f(V(L_n)) = \{1, 2, 3, \dots, 2n\}$; for $1 \leq i \leq n-m$,

$$\begin{aligned} f(x_i) + f(y_i) &= f(x_{m+i}) + f(y_{m+i}) - 2, \\ f(x_i y_i) + f(x_{m+i-1} y_{m+i-1}) &= f(x_{i+1} y_{i+1}) + f(x_{m+i} y_{m+i}) + 2; \end{aligned}$$

for $1 \leq i \leq n-2$,

$$f(x_i x_{i+1}) + f(y_i y_{i+1}) = f(x_{i+1} x_{i+2}) + f(y_{i+1} y_{i+2}).$$

For $1 \leq i \leq n-m+1$, let $C_{2m}^{(i)}$ be the subcycle of L_n with

$$V(C_{2m}^{(i)}) = \{x_j, y_j | i \leq j \leq i+m-1\}$$

and

$$E(C_{2m}^{(i)}) = \{x_j x_{j+1}, y_j y_{j+1} | i \leq j \leq i+m-2\} \cup \{x_i y_i, x_{i+m-1} y_{i+m-1}\}.$$

It is easy to verify that $V(C_{2m}^{(i)}) \cap V(C_{2m}^{(i+1)}) = \{x_j, y_j | i+1 \leq j \leq i+m-1\}$ and $E(C_{2m}^{(i)}) \cap E(C_{2m}^{(i+1)}) = \{x_j x_{j+1}, y_j y_{j+1} | i+1 \leq j \leq i+m-2\}$.

By using these facts, for $1 \leq i \leq n-m$, we obtain

$$\begin{aligned} \sum f(C_{2m}^{(i)}) &= \sum_{j=i}^{i+m-1} [f(x_j) + f(y_j)] + \sum_{j=i}^{i+m-2} [f(x_j x_{j+1}) + f(y_j y_{j+1})] + \\ &\quad f(x_i y_i) + f(x_{i+m-1} y_{i+m-1}) \\ &= \sum_{j=i+1}^{i+m} [f(x_j) + f(y_j)] + \sum_{j=i+1}^{i+m-1} [f(x_j x_{j+1}) + f(y_j y_{j+1})] + \\ &\quad f(x_{i+1} y_{i+1}) + f(x_{i+m} y_{i+m}) \\ &= \sum f(C_{2m}^{(i+1)}). \end{aligned}$$

So, for $1 \leq i \leq n-m+1$, $\sum f(C_{2m}^{(i)})$ is constant. Hence, f is a C_{2m} -supermagic labeling of L_n . \square




Figure 3: (a). a C_6 -supermagic labeling of L_6 , (b). a C_8 -supermagic labeling of L_6

In Figure 3 we show a C_6 -supermagic labeling and a C_8 -supermagic labeling of L_6 as defined in the proof of Theorem 3.

Next, we consider a L_m -supermagic labeling of L_n . Notice that, L_n is $L_2 \cong C_4$ -supermagic and L_n is trivially L_n -supermagic. So, in the following theorem, we consider a L_m -supermagic labeling of L_n for any integer $3 \leq m \leq n - 1$.

Theorem 4. *Let $n \geq 4$ be a positive integer. Then the ladder L_n is L_m -supermagic for every integer $3 \leq m \leq n - 1$.*

Proof. For proving this theorem, we define the ladder L_n as a graph with $V(L_n) = \{x_i, y_i | 1 \leq i \leq n\}$ and $E(L_n) = \{x_i x_{i+1}, y_i y_{i+1} | 1 \leq i \leq n - 1\} \cup \{x_i y_{n+1-i} | 1 \leq i \leq n\}$.

Define a total labeling of L_n in the following way.

- For $1 \leq i \leq n$, label x_i as in the proof of Theorem 3.
- For $1 \leq i \leq n$, label y_i with $n + (\text{the label of } x_{n+1-i})$.
- For $1 \leq i \leq n - 1$, label $x_i x_{i+1}$ with $2n + i$.
- For $1 \leq i \leq n$, label $x_i y_{n+1-i}$ with $5n - 1 - i$.

For labeling $y_i y_{i+1}$, we consider two following cases. First, let $q = n - 2$.

Case 1: $q \equiv 0 \pmod{m - 1}$

- Label $y_1 y_2, y_m y_{m+1}, y_{2m-1} y_{2m}, y_{3m-2} y_{3m-1}, y_{4m-3} y_{4m-2}, \dots, y_{n-m} y_{n-m+1}, y_{n-1} y_n$ with $3n, 3n + 1, 3n + 2, 3n + 3, 3n + 4, 3n + 5, \dots, 3n + (\frac{q}{m-1} - 1), 3n + (\frac{q}{m-1})$, respectively.

- For $1 \leq k \leq m-2$, label $y_{1+k}y_{2+k}, y_{m+k}y_{m+k+1}, y_{2m+k-1}y_{2m+k}, y_{3m+k-2}y_{3m+k-1}, y_{4m+k-3}y_{4m+k-2}, \dots, y_{n-m+k}y_{n-m+k+1}$ with $3n + k(\frac{q}{m-1}) + 1, 3n + k(\frac{q}{m-1}) + 2, 3n + k(\frac{q}{m-1}) + 3, 3n + k(\frac{q}{m-1}) + 4, 3n + k(\frac{q}{m-1}) + 5, \dots, 3n + (k+1)(\frac{q}{m-1})$, respectively.

Case 2: $q \equiv t \pmod{m-1}$, where $t = 1, 2, 3, \dots, m-2$

- Label $y_1y_2, y_my_{m+1}, y_{2m-1}y_{2m}, y_{3m-2}y_{3m-1}, y_{4m-3}y_{4m-2}, \dots, y_{q-t-m+2}y_{q-t-m+3}, y_{q-t+1}y_{q-t+2}$ with $3n, 3n+1, 3n+2, 3n+3, 3n+4, 3n+5, \dots, 3n + (\frac{q-t}{m-1} - 1), 3n + (\frac{q-t}{m-1})$, respectively.
- Label $y_\alpha y_{\alpha+1}, y_{\alpha+1}y_{\alpha+2}, y_{\alpha+2}y_{\alpha+3}, \dots, y_{\alpha+t-1}y_{\alpha+t}$ with $3n+2(\frac{q-t}{m-1})+1, 3n+3(\frac{q-t}{m-1})+2, 3n+4(\frac{q-t}{m-1})+3, \dots, 3n+(t+1)(\frac{q-t}{m-1})+t$, respectively, where $\alpha = q-t+2$.
- For $1 \leq k \leq t+1$, label $y_{k+1}y_{k+2}, y_{m+k}y_{m+k+1}, y_{2m+k-1}y_{2m+k}, y_{3m+k-2}y_{3m+k-1}, y_{4m+k-3}y_{4m+k-2}, \dots, y_{q-t-m+k+2}y_{q-t-m+k+3}$, with $\beta_1^k, \beta_1^k + 1, \beta_1^k + 2, \beta_1^k + 3, \beta_1^k + 4, \dots, \beta_1^k + (\frac{q-t}{m-1} - 1)$, respectively, where $\beta_1^k = k(\frac{q-t}{m-1} + 1) + 3n$.
- For $t+2 \leq k \leq m-2$, label $y_{k+1}y_{k+2}, y_{m+k}y_{m+k+1}, y_{2m+k-1}y_{2m+k}, y_{3m+k-2}y_{3m+k-1}, y_{4m+k-3}y_{4m+k-2}, \dots, y_{q-t-m+k+2}y_{q-t-m+k+3}$, with $\beta_2^k, \beta_2^k + 1, \beta_2^k + 2, \beta_2^k + 3, \beta_2^k + 4, \dots, \beta_2^k + (\frac{q-t}{m-1} - 1)$, respectively, where $\beta_2^k = k(\frac{q-t}{m-1}) + 3n + t + 1$.

Let us denote the labeling defined above by g . For $1 \leq i \leq n-m+1$, it can be checked that

$$\begin{aligned} \sum_{j=i}^{i+m-1} [g(x_j) + g(y_{n+1-j})] &= -2 + \sum_{j=i+1}^{i+m} [g(x_j) + g(y_{n+1-j})], \\ \sum_{j=i}^{i+m-2} g(x_j x_{j+1}) &= 1 - m + \sum_{j=i+1}^{i+m-1} g(x_j x_{j+1}), \\ \sum_{j=i}^{i+m-2} g(y_{n+1-j} y_{n-j}) &= 1 + \sum_{j=i+1}^{i+m-1} g(y_{n+1-j} y_{n-j}), \end{aligned}$$

and

$$\sum_{j=i}^{i+m-1} g(x_j y_{n+1-j}) = m + \sum_{j=i+1}^{i+m} g(x_j y_{n+1-j}).$$

For $1 \leq i \leq n-m+1$, let $L_m^{(i)}$ be the subladder of L_n with $V(L_m^{(i)}) = \{x_j, y_{n+1-j} | i \leq j \leq m+i-1\}$ and $E(L_m^{(i)}) = \{x_j x_{j+1}, y_{n+1-j} y_{n-j} | i \leq j \leq m+i-2\} \cup \{x_j y_{n+1-j} | i \leq j \leq m+i-1\}$.

In a similar way as in the proof of Theorem 3, for $1 \leq i \leq n - m$, it is easy to verify that $\sum g(L_m^{(i)}) = \sum g(L_m^{(i+1)})$.

So, $\sum g(L_m^{(i)})$ is constant for all possible values of i . Hence, L_n is L_m -supermagic for every integer $3 \leq m \leq n - 1$. \square

An example of the labeling obtained in the above proof is showed in Figure 4.




Figure 4: an L_3 -supermagic labeling of L_9

Acknowledgement

The authors are thankful to the anonymous referee for some comments that helped to improve the presentation of the manuscript. The first and the second authors were funded by “Hibah Kompetensi” from the Directorate General of Higher Education of Indonesia. The third author was supported by “Insentif Riset Dasar” from the State Ministry of Research and Technology of Indonesia.

References

- [1] M. Bača, *On magic labelings of convex polytopes*, Annals of Discrete Math. **51** (1992), 13-16.
- [2] H. Enomoto, A. Llado, T. Nakamigawa, and G. Ringel, *Super edge magic graphs*, SUT J. Math. **34** (1998), 105-109.
- [3] Y. Fukuchi, *A recursive theorem for super edge-magic labelings of trees*, SUT J. Math. **36**:2 (2000), 279-285.
- [4] J.A. Gallian, *A dynamic survey of graph labelings*, Electron. J. Combin. **16** (2009), # DS6.
- [5] A. Gutiérrez and A. Lladó, *Magic coverings*, J. Combin. Math. Combin. Comput. **55** (2005), 43-56.
- [6] A. Kotzig and A. Rosa, *Magic valuation of finite graphs*, Canad. Math. Bull. **13**:4 (1970), 451-461.
- [7] K.W. Lih, *On magic and consecutive labelings of plane graphs*, Utilitas Math. **24** (1983), 165-197.

- [8] A. Lladó and J. Moragas, *Cycle-magic graphs*, Discrete Math. **307**:23 (2007), 2925-2933.
- [9] T. K. Maryati, E.T. Baskoro, and A.N.M. Salman, *P_h -supermagic labelings of some trees*, J. Combin. Math. Combin. Comput. **65** (2008), 197-204.
- [10] T. K. Maryati, A.N.M. Salman, E.T. Baskoro, J. Ryan, and M. Miller, *On H -supermagic labelings for certain shackles and amalgamations of a connected graph*, Utilitas Math. to appear.
- [11] A.A.G. Ngurah, E.T. Baskoro, R. Simanjuntak, and S. Uttunggadewa, *On super edge-magic strength and deficiency of graphs*, Lecture Notes in Comput. Sci. **4535** (2008), 144-154.
- [12] A.A.G. Ngurah, E.T. Baskoro, and I. Tomescu, *Magic graphs with pendant edges*, Ars Combin. to appear.
- [13] A.A.G. Ngurah, A.N.M. Salman, and L. Susilowati, *H -supermagic labelings of graphs*, Discrete Math. **310**:8 (2010), 1293-1300.
- [14] A.A.G. Ngurah, R. Simanjuntak, and E.T. Baskoro, *On (super) edge-magic total labeling of a subdivision of $K_{1,3}$* , SUT J. Math. **43**:2 (2007), 127-136.
- [15] A.N.M. Salman, A.A.G. Ngurah, and N. Izzati, *On (super) edge-magic total labeling of subdivision of stars S_n* , Utilitas Math. to appear.

A.A.G. Ngurah

Department of Civil Engineering, Universitas Merdeka Malang

Jalan Taman Agung No. 1 Malang 65146, Indonesia

E-mail: ngurahram67@yahoo.com

A.N.M. Salman

Combinatorial Mathematics Research Group, Faculty of Mathematics and Natural Sciences

(FMIPA), Institut Teknologi Bandung (ITB)

Jalan Ganesa No. 10 Bandung 40132, Indonesia

E-mail: msalman@math.itb.ac.id

I W. Sudarsana

Combinatorial and Applied Mathematics Research Group, Department of Mathematics,

Faculty of Mathematics and Natural Sciences (FMIPA), Tadulako University (UNTAD)

Jalan Soekarno-Hatta Km. 8 Palu 94118, Indonesia

E-mail: sudarsanaiwayan@yahoo.co.id