


Curriculum Vitae

Hideshi Yokoyama, Ph. D.

Professor



Current Address:

Laboratory of Physical Biochemistry, Faculty of Pharmaceutical Sciences,

Tokyo University of Science

2641 Yamazaki, Noda, Chiba 278-8510, Japan

TEL&FAX: 81-4-7121-3663

Email: yokoyama(add @ here)rs.tus.ac.jp

URL: http://www.rs.tus.ac.jp/yokoyama/English/index_e.html

Personal Information:

Name, Family name: Yokoyama

Forenames: Hideshi

Sex: Male

Nationality: Japanese

Date of birth: March 1974

Research and professional experience:

- | | |
|------------------|--|
| 2001 – 2002 | Predocctoral Research Fellow (DC2) of Japan Society for the Promotion of Science, Japan |
| 2002 – 2006 | Technical Staff, Biological Information Research Center, National Institute of Advanced Industrial Science and Technology, Japan |
| 2006 – 2011 | Research Assistant Professor, School of Pharmaceutical Sciences, University of Shizuoka, Japan |
| 2010.8 – 2010.11 | Visiting Researcher, Institut de Génétique et Microbiologie, Université Paris-Sud (Professor Patrick Forterre), France |
| 2011 – 2016 | Assistant Professor, School of Pharmaceutical Sciences, University of Shizuoka, Japan |
| 2016 – 2023 | Associate Professor, Faculty of Pharmaceutical Sciences, Tokyo University of Science, Japan |
| 2023 – present | Professor, Faculty of Pharmaceutical Sciences, Tokyo University of Science, Japan |

Education:

1993 – 1997 Faculty of Pharmaceutical Sciences,
University of Tokyo, Japan
1997 – 2002 Graduate School of Pharmaceutical Sciences,
University of Tokyo, Japan
2002 Obtained Ph. D.

Award:

Award for Young Scientists in Tokai Branch of Pharmaceutical Society of Japan, Japan
(2009)

Research interest:

Structural biology of membrane signalling proteins or drug-target proteins using X-ray crystallography

Original Paper:

(*Corresponding author, #Contributed equally)

56. Hydrophobic hydration of analgesics and diltiazem complexes explored by electrochemical impedance spectroscopy and diffusion-ordered spectroscopy.

Koga, R., Kinoshita, T., Ishiguro, M., Fujita, M., Chatani, H., Kataoka, H., Yokoyama, H., Hanawa, T., Shitanda, I. & *Goto, S.

Phys. Chem. Chem. Phys. **28**, 1159-1172 (2026).

55. Comparative study on the effects of the inclusion complexes of non-steroidal anti-inflammatory drugs with 2-hydroxypropyl- β -cyclodextrins on dissociation rates and supersaturation.

Oshite, Y., Wada-Hirai, A., Ichii, R., Kuroda, C., Hasegawa, K., Hiroshige, R., Yokoyama, H., Tsuchida, T. & *Goto, S.

RSC Pharm. **1**, 80-97 (2024).

54. Quantification of crystallinity during indomethacin crystalline transformation from α - to γ -polymorphic forms and of the thermodynamic contribution to dissolution in aqueous buffer and solutions of solubilizer.

Hasegawa, K., *Goto, S., Kataoka, H., Chatani, H., Kinoshita, T., Yokoyama, H. & Tsuchida, T.

RSC Adv. **14**, 4129-4141 (2024).

53. Wnt activation disturbs cell competition and causes diffuse invasion of transformed cells through NF- κ B-MMP21 pathway.

Nakai, K., Lin, H., Yamano, S., Tanaka, S., Kitamoto, S., Saitoh, H., Sakuma, K., Kurauchi, J., Akter, E., Konno, M., Ishibashi, K., Kamata, R., Ohashi, A., Koseki, J., Takahashi, H., Yokoyama, H., Shiraki, Y., Enomoto, A., Abe, S., Hayakawa, Y., Ushiku, T., Mutoh, M., Fujita, Y. & Kon, S.

Nat. Commun. **14**, 7048 (2023).

52. Using singular value decomposition to analyze drug/ β -cyclodextrin mixtures: insights from X-ray powder diffraction patterns.

Hasegawa, K., *Goto, S., Tsunoda, C., Kuroda, C., Okumura, Y., Hiroshige, R., Wada-Hirai, A., Shimizu, S., Yokoyama, H. & Tsuchida, T.

Phys. Chem. Chem. Phys. **25**, 29266-29282 (2023).

51. Effect of cyclodextrin complex formation on solubility changes of each drug due to intermolecular interactions between acidic NSAIDs and basic H₂ blockers.

Tsunoda, C., Hasegawa, K., Hiroshige, R., Kasai, T., Yokoyama, H. & *Goto, S.

Mol. Pharm **20**, 5032-5042 (2023).

50. Effects of local anesthetics on liposomal membranes determined by their inhibitory activity of lipid peroxidation.

Horizumi, Y., *Goto, S., Takatsuka, M. & Yokoyama, H.

Mol. Pharm **20**, 2911-2918 (2023).

49. Higher-order structure formation using refined monomer structures of lipid raft markers, Stomatin, Prohibitin, Flotillin, and HflK/C-related proteins.

Yokoyama, H. & *Matsui, I.

FEBS Open Bio **13**, 926-937 (2023).

48. Crystal structure of the motor domain of centromere-associated protein E in complex with a non-hydrolysable ATP analogue.

Shibuya, A., Suzuki, A., Ogo, N., Sawada, J., Asai, A. & *Yokoyama, H.

FEBS Lett. **597**, 1138-1148 (2023).

47. Optimization of the stability constants of the ternary system of diclofenac/famotidine/ β -cyclodextrin by nonlinear least-squares method using theoretical equations.

Tsunoda, C., *Goto, S., Hiroshige, R., Kasai, T., Okumura, Y. & Yokoyama, H.

Int. J. Pharm. **638**, 122913 (2023).

46. Interaction mode of hydroxypropyl- β -cyclodextrin with vaccine adjuvant components Tween 80 and Triton X-100 revealed by fluorescence increasing-quenching analysis.

Kurosawa, Y., *Goto, S., Mitsuya, K., Otsuka, Y. & Yokoyama, H.

Phys. Chem. Chem. Phys. **25**, 6203-6213 (2023).

45. Structural and mutational studies suggest key residues to determine whether stomatin SPFH domains form dimers or trimers.

Komatsu, T., Matsui, I. & *Yokoyama, H.

Biochem. Biophys. Rep. **32**, 101384 (2022).

44. Trajectory of the spectral/structural rearrangements for photo-oxidative reaction of neat ketoprofen and its cyclodextrin complex.

Hiroshige, R., *Goto, S., Tsunoda, C., Ichii, R., Shimizu, S., Otsuka, Y., Makino, K., Takahashi, H. & Yokoyama, H.

J. Incl. Phenom. Macrocycl. Chem. **102**, 791-800 (2022).

43. Fluorescence changes of dyes/NSAIDs adsorbed on fluorocarbon polymers.

Mitsuya, K., *Goto, S., Kurosawa, Y., Yokoyama, H. & Hanawa, T.

Mater. Chem. Phys. **290**, 126552 (2022).

42. Structural basis for peptide recognition by archaeal oligopeptide permease A.

*Yokoyama, H., Kamei, N., Konishi, K., Hara, K., Ishikawa, Y., Matsui, I., Forterre, P. & Hashimoto, H.

Proteins: Struct Funct Bioinf. **90**, 1434-1442 (2022).

41. Crystal structures of human glyoxalase I and its complex with TLSC702 reveal inhibitor binding mode and substrate preference.

Usami, M., Ando, K., Shibuya, A., Takasawa, R. & *Yokoyama, H.

FEBS Lett. **596**, 1458-1467 (2022).

40. Preparation, crystallization, and X-ray data collection of archaeal oligopeptide permease A.

*Yokoyama, H., Kamei, N., Konishi, K., Hara, K., Ishikawa, Y., Matsui, I., Forterre, P. & Hashimoto, H.

Crystallogr. Rep. **66**, 1300-1305 (2021).

39. Structure and comparison of the motor domain of centromere-associated protein E.

Shibuya, A., Ogo, N., Sawada, J., Asai, A. & *Yokoyama, H.

Acta Crystallog. **D77**, 280-287 (2021).

38. Inactive dimeric structure of the protease domain of stomatin operon partner protein.

*Yokoyama, H., Suzuki, K., Hara, K., Matsui, I. & Hashimoto, H.

Acta Crystallog. **D76**, 515-520 (2020).

37. The lipid raft markers stomatin, prohibitin, flotillin, and HflK/C (SPFH)-domain proteins form an operon with NfeD proteins and function with apolar polyisoprenoid lipids.

Yokoyama, H. & *Matsui, I.

Crit. Rev. Microbiol. **46**, 38-48 (2020).

36. Structural and biochemical basis of the formation of isoaspartate in the complementarity-determining region of antibody 64M-5 Fab.

Yokoyama, H., Mizutani, R., Noguchi, S. & *Hayashida, N.

Sci. Rep. **9**, 18494 (2019).

35. Structures of the antibody 64M-5 Fab and its complex with dT(6-4)T indicate induced-fit and high-affinity mechanisms.

Yokoyama, H., Mizutani, R., Noguchi, S. & *Hayashida, N.

Acta Crystallog. **F75**, 80-88 (2019).

34. Molecular modeling and simulation of human stomatin and predictions for its membrane association.

Kondo, Y., *Yokoyama, H., Matsui, I. & Miyazaki, S.

J. Data Mining Genomics Proteomics, **9**, 216 (2018).

33. Structural and thermodynamic basis of the enhanced interaction between kinesin spindle protein Eg5 and STLC-type inhibitors.

*[#]Yokoyama, H., [#]Sawada, J., Sato, K., Ogo, N., Kamei, N., Ishikawa, Y., Hara, K., *Asai, A. & Hashimoto, H.

ACS Omega, **3**, 12284-12294 (2018).

32. Structure of proliferating cell nuclear antigen (PCNA) bound to an APIM peptide reveals the universality of PCNA interaction.

Hara, K., Uchida, M., Tagata, R., Yokoyama, H., Ishikawa, Y., Hishiki, A. & *Hashimoto, H.

Acta Crystallog. **F74**, 214-221 (2018).

31. Dynamic feature of mitotic arrest deficient 2-like protein 2 (MAD2L2) and structural basis for its interaction with chromosome alignment-maintaining phosphoprotein (CAMP).

Hara, K., Taharazako, S., Ikeda, M., Fujita, H., Mikami, Y., Kikuchi, S., Hishiki, A., Yokoyama, H., Ishikawa, Y., Kanno, SI., Tanaka, K. & *Hashimoto, H.

J. Biol. Chem. **292**, 17658-17667 (2017).

30. Role of the mobility of antigen binding site in high affinity antibody elucidated by surface plasmon resonance.

Fukuda, N., Suwa, Y., Uchida, M., Kobashigawa, Y., Yokoyama, H. & *Morioka, H.
J. Biochem. **161**, 37-43 (2017)

29. Structure of a novel DNA-binding domain of helicase-like transcription factor (HLTF) and its functional implication in DNA damage tolerance.

Hishiki, A., Hara, K., Ikegaya, Y., Yokoyama, H., Shimizu, T., Sato, M. & *Hashimoto, H.
J. Biol. Chem. **290**, 13215-13223 (2015).

28. Crystallographic study of a novel DNA-binding domain of human HLTF involved in the template-switching pathway to avoid the replication arrest caused by DNA damage.

Ikegaya, Y., Hara, K., Hishiki, A., Yokoyama, H. & *Hashimoto, H.
Acta Crystallog. **F71**, 668-670 (2015).

27. Structural basis of new allosteric inhibition in kinesin spindle protein Eg5.

*Yokoyama, H., #Sawada, J., Katoh, S., Matsuno, K., Ogo, N., Ishikawa, Y., Hashimoto, H., Fujii, S. & *Asai, A.
ACS Chem. Biol. **10**, 1128-1136 (2015).

26. Structural biology of DNA (6-4) photoproducts formed by ultraviolet radiation and interactions with their binding proteins.

*Yokoyama, H. & Mizutani, R.
Int. J. Mol. Sci. **15** (11), 20321-20338 (2014).

25. Crystal structure of the stomatin operon partner protein from *Pyrococcus horikoshii* indicates the formation of a multimeric assembly.

*Yokoyama, H. & Matsui, I.
FEBS Open Bio **4**, 804-812 (2014).

24. Structures and metal-binding properties of *Helicobacter pylori* neutrophil-activating protein with a di-nuclear ferroxidase center.

*Yokoyama, H. & Fujii, S.
Biomolecules **4** (3), 600-615 (2014).

23. Structural and biochemical analysis of a thermostable membrane-bound stomatin-specific protease.

*Yokoyama, H., Kobayashi, D., Takizawa, N., Fujii, S. & Matsui, I.
J. Synchrotron Rad. **20**, 933-937 (2013).

22. Domain Structures and Inter-Domain Interactions Defining the Holoenzyme Architecture of Archaeal D-Family DNA Polymerase.

*Matsui, I., Matsui, E., Yamasaki, K. & Yokoyama, H.
Life **3** (3), 375-385 (2013).

21. Clustering of OB-fold domains of the partner protease complexed with trimeric stomatin from Thermococcales.

Yokoyama, H., Matsui, E., Hiramoto, K., Forterre, P. & *Matsui, I.
Biochimie **95**, 1494-1501 (2013).

20. Structure of a double-stranded DNA (6-4) photoproduct in complex with the 64M-5 antibody Fab.

*Yokoyama, H., Mizutani, R. & Satow, Y.
Acta Crystallog. **D69**, 504-512 (2013).

19. Crystal structure of *Helicobacter pylori* neutrophil-activating protein with a di-nuclear ferroxidase center in a zinc or cadmium-bound form.

*Yokoyama, H., Tsuruta, O., Akao, N. & Fujii, S.
Biochem. Biophys. Res. Commun. **422**, 745-750 (2012).

18. Crystal structure of a membrane stomatin-specific protease in complex with a substrate peptide.

*Yokoyama, H., Takizawa, N., Kobayashi, D., Matsui, I. & Fujii, S.
Biochemistry **51**, 3872-3880 (2012).

17. Structure of the DNA (6-4) photoproduct dTT(6-4)TT in complex with the 64M-2 antibody Fab fragment implies increased antibody-binding affinity by the flanking nucleotides.

*Yokoyama, H., Mizutani, R., Satow, Y., Sato, K., Komatsu, Y., Ohtsuka, E. & Nikaido, O.
Acta Crystallog. **D68**, 232-238 (2012).

16. A new crystal lattice structure of *Helicobacter pylori* neutrophil-activating protein (HP-NAP).

Tsuruta, O., *Yokoyama, H. & Fujii, S.
Acta Crystallog. **F68**, 134-140 (2012).

15. Novel structure of an N-terminal domain that is crucial for the dimeric assembly and DNA-binding of an archaeal DNA polymerase D large subunit from *Pyrococcus horikoshii*.

*Matsui, I., Urushibata, Y., Shen, Y., Matsui, E. & Yokoyama, H.
FEBS Lett. **585**, 452-458 (2011).

14. Three-dimensional structure of membrane protein stomatin and function of stomatin-specific protease.

*Yokoyama, H.

Yakugaku Zasshi **130(10)**, 1289-1293 (2010).

13. Binding of sulphatide to recombinant haemagglutinin of influenza A virus produced by a baculovirus protein expression system.

Takahashi, T., Satoh, H., Takaguchi, M., Takafuji, S., Yokoyama, H., Fujii, S. & *Suzuki, T.
J. Biochem. **147**, 459-462 (2010).

12. Unusual thermal disassembly of the SPFH domain oligomer from *Pyrococcus horikoshii*.

Kuwahara, Y., Unzai, S., Nagata, T., Hiroaki, Y., Yokoyama, H., Matsui, I., Ikegami, T., Fujiyoshi, Y. & *Hiroaki, H.

Biophys. J. **97**, 2034-2043 (2009).

11. The solution structure of the C-terminal domain of NfeD reveals a novel membrane-anchored OB-fold.

Kuwahara, Y., Ohno, A., Morii, T., Yokoyama, H., Matsui, I., Tochio, H., Shirakawa, M. & *Hiroaki, H.

Protein Sci. **17**, 1915-1924 (2008).

10. Novel dimer structure of a membrane-bound protease with a catalytic Ser-Lys dyad and its linkage to stomatin.

Yokoyama, H., Hamamatsu, S., Fujii, S. & *Matsui, I.

J. Synchrotron Rad. **15**, 254-257 (2008).

9. Crystal structure of a core domain of stomatin from *Pyrococcus horikoshii* illustrates a novel trimeric and coiled-coil fold.

Yokoyama, H., Fujii, S. & *Matsui, I.

J. Mol. Biol. **376**, 868-878 (2008).

8. Molecular structure of a novel membrane protease specific for a stomatin homolog from the hyperthermophilic archaeon *Pyrococcus horikoshii*.

Yokoyama, H., Matsui, E., Akiba, T., Harata, K. & *Matsui, I.

J. Mol. Biol. **358**, 1152-1164 (2006).

7. A novel thermostable membrane protease forming an operon with a stomatin homolog from the hyperthermophilic archaeobacterium *Pyrococcus horikoshii*.

Yokoyama, H. & *Matsui, I.

J. Biol. Chem. **280**, 6588-6594 (2005).

6. Aromatic residues located close to the active center are essential for the catalytic reaction of flap endonuclease-1 from hyperthermophilic archaeon *Pyrococcus horikoshii*.

Matsui, E., Abe, J., Yokoyama, H. & *Matsui, I.

J. Biol. Chem. **279**, 16687-16696 (2004).

5. A 21-amino acid peptide from the cysteine cluster II of the family D DNA polymerase from *Pyrococcus horikoshii* stimulates its nuclease activity which is Mre11-like and prefers manganese ion as the cofactor.

Shen, Y., Tang, X.-F., Yokoyama, H., Matsui, E. & *Matsui, I.

Nucleic Acids Res. **32**, 158-168 (2004).

4. Distinct domain functions regulating de novo DNA synthesis of thermostable DNA primase from hyperthermophile *Pyrococcus horikoshii*.

Matsui, E., Nishio, M., Yokoyama, H., Harata, K., Darnis, S. & *Matsui, I.

Biochemistry **42**, 14968-14976 (2003).

3. Complex formation of double-stranded DNA (6-4) photoproducts and anti-(6-4) photoproduct antibody Fabs.

Yokoyama, H., Mizutani, R. & *Satow, Y.

Nucleic Acids Symp. Ser. **44**, 151-152 (2000).

2. Crystal structure of the 64M-2 antibody Fab fragment in complex with a DNA dT(6-4)T photoproduct formed by ultraviolet radiation.

Yokoyama, H., Mizutani, R., *Satow, Y., Komatsu, Y., Ohtsuka, E. & Nikaido, O.

J. Mol. Biol. **299**, 711-723 (2000).

1. Crystal structures of the 64M-2 and 64M-3 antibody Fabs complexed with DNA (6-4) photoproducts.

Yokoyama, H., Mizutani, R., *Satow, Y., Komatsu, Y., Ohtsuka, E. & Nikaido, O.

Nucleic Acids Symp. Ser. **42**, 267-268 (1999).